
Evaluation of an AI Primary Eye Care Agent

Physician-Validated Multi-LLM-as-a-Judge Assessment Across Frontier and Local Models in Low-Resource Settings

MSc in Health Data Science

Matriculation Number - 250009792



University of
St Andrews

Supervised by Dr Andrew Blaikie, Dr Craig Myles and William J. Williams

August 2026

Declaration

I hereby certify that this project, which is approximately 13,400 words in length, has been composed by me, that it is the record of work carried out by me, and that it has not been submitted in any previous application for a degree.

This project was conducted by me at the University of St Andrews from June 2026 to August 2026 towards fulfilment of the requirements of the University of St Andrews for the degree of MSc in Health Data Science under the supervision of Dr Andrew Blaikie, Dr Craig Myles and William J. Williams.

Matriculation Number : 250009792

Dated: 10/08/2026



Acknowledgement

I would like to thank my supervisor, Andrew Blaikie, and my co-advisors, Craig Myles and William J. Williams, for their generous time, encouraging feedback and continuous support throughout this project. Andrew's encouragement gave me the confidence to grow beyond what I thought I could do. His commitment to global health continues to inspire me, and I am fortunate to have learned under it. Craig's insights laid the foundations for a robust methodology and a rigorous engagement with the literature. William's thoughtful questions consistently encouraged me to probe my own assumptions and sustained both my curiosity and desire to read more widely on the subject. I am equally grateful to Henok S. Hailemichael, whose dedicated time and thoughtful interventions kept the study on track and clinically relevant.

Abstract

An estimated 1.1 billion people live with untreated vision impairment, most of them in low- and middle-income countries where eye specialists are scarce. One response to that scarcity is Alan, an open-source clinical scaffold that can guide a frontline eye consultation on any large language model (LLM). Frontier models are costly and cloud-bound, while small open-weight models can run inside the clinic itself, yet no study had measured what is lost when the model beneath the scaffold changes. This study set out to quantify exactly that.

Alan was evaluated on seven models, two frontier and five open-weight, each with extended reasoning ("thinking") enabled and disabled. Every configuration worked through 48 ophthalmic cases presented by two simulated health-worker personas, yielding 1,344 consultations in total. Local models ran on the University of St Andrews Hypatia HPC cluster with energy measured throughout. A blinded panel of three LLM judges then scored every dialogue on a five-criterion weighted rubric, anchored by two human ophthalmologists.

The results were closer than expected. The best frontier configuration scored 84.2 out of 100, while the best open-weight model reached 81.8, a gap of just 2.4 points (95% CI 0.3 to 4.8). Below this leading group, however, quality fell steeply to a floor of 51.4. The LLM judges reproduced the ophthalmologists' model ranking, but their pass rate was substantially lower (54.1% vs 79.8%), which supports their use for comparative ranking rather than as a direct estimate of clinical acceptability. In deployment terms, latency rather than cost emerged as the principal constraint: the highest-scoring local configuration took 64.8 seconds per reply, whereas the fastest deployable one answered in under six seconds while scoring 77.0.

Taken together, these findings show that Alan produced different clinical behaviour on different hosts. This reinforces the case for regulators to treat the scaffold-model combination as one inseparable product, and for health systems to demand evidence from hardware and configurations their clinics can realistically operate.

Glossary

Agent	Software that can act on its own; taking in information, making decisions and carrying out steps toward a goal.
Context	How much text a model can hold in mind at once, measured in tokens.
Host Model	The underlying model that a scaffold or prompt runs on.
Open-source	Software or models whose code and design are published freely, so anyone can inspect, change, and reuse them.
Open-weight	A model whose trained parameters (weights) are released publicly, so it can be run locally. Though its training data or code may stay private.
Persona	A role given to a model by instruction (for example, a patient or an eye specialist) to shape how it behaves.
Preset	A fixed set of model settings used so that every run behaves the same way.
Prompt	The text or instruction given to a model to obtain a response.
Scaffold	An external code, prompt, memory systems and tool integrations built around a model to guide its behaviour and execute complex tasks (here, ‘Alan’).
Temperature	A setting that controls randomness. Near 0 the model is predictable; higher values make it more varied.
Token	The basic chunk of text a model reads and writes.

Table of Contents

1. Introduction 1

1.1 Background and motivation 1

1.2 Alan: an open-source clinical scaffold 1

1.3 The problem: model scale and the deployment gap 2

1.4 Research questions 2

1.5 Significance of this study 3

1.6 Scope of the study 3

2. Literature Review 5

2.1 Task shifting and the Arclight 5

2.2 The barriers to global digital health 5

2.3 RAG and commercial evidence tools as alternatives 6

2.4 The scaffold-model pair and instruction following 7

2.5 LLMs in eye care and their evaluation 8

2.6 LLM-as-a-judge, rubrics and the reasoning 9

2.7 Chain-of-thought reasoning as diagnostic evidence 10

2.8 Regulation and governance of clinical LLMs 11

2.9 The gap this study occupies 12

3. Methodology 14

3.1 Methodology overview 14

3.2 Simulation as the evaluation method 20

3.3 Evaluation rubric 20
3.4 Case bank 22
3.5 The scaffold 23
3.6 The simulated health worker 23
3.7 The LLM judge panel 23
3.8 Human expert validation 24
3.9 Dialogue orchestration and environment 25
3.10 Blinding and bias controls 27
3.11 Statistical Analysis 27
3.12 Energy footprint and running cost estimation 28
3.13 Reproducibility and data availability 28
3.14 Ethics 28
4. Results 29
4.1 Dialogues 29
4.2 Score data 31
4.3 Leaderboard 33
4.4 Frontier to local gap 36
4.5 Thinking on vs thinking off 38
4.6 Adherence against accuracy 39
4.7 The health worker personas 41
4.8 The LLM judge panel as an instrument 42

4.9 Physician validation 43

4.10 Robustness of the ranking 44

4.11 Deployment measures: latency, energy and cost 45

4.12 Sensitivity check 48

5. Discussion 49

5.1 What the study found 49

5.2 Why the gap is smaller than the literature predicted 49

5.3 Following the rules is not the same as being right 50

5.4 Lowest scores, endemic diseases: a training-data problem? 51

5.5 LLM judge panel performance, human ceilings and criterion reliability 52

5.6 Thinking, latency, and what it costs 54

5.7 Two owners of one product and what this study cannot say 55

6. Conclusion 56

References 58

Appendix 66

Appendix I: Ethical Approval 66

Appendix II: Evaluation Rubric 67

Appendix III: Health Worker Persona System Prompts 71

Appendix IV: Expert Judge Persona System Prompts 76

Appendix V: Snapshot of Alan, the scaffold 79

Appendix VI: LMIC medication and instruments reference adopted from WHO package of eyecare interventions [106] 80

Appendix VII: Dialogue completion by preset 81

Appendix VIII: Paired effect of thinking on 82

Appendix IX: Clinical accuracy against scaffold adherence 83

Appendix X: Weighted composite by health worker persona 84

List of Tables

Table 1: The evaluation rubric, its weights and the reason each criterion was included. 20

Table 2: Example record from the case set, case 01. 22

Table 3: Every model used, with source, serving backend and settings 26

Table 4: Leaderboard of all fourteen presets, with case-clustered bootstrap 95 percent intervals on the weighted composite. Accept % is the share of each preset's 576 judgements that cleared the gate. 33

Table 5: Shortfall against the leading preset (GPT-5.4 with thinking on), paired case by case, with case-clustered bootstrap intervals. 36

Table 6: LLM judges against the pooled physician mean, 134 dialogues. Shift is judge minus physicians on the 25-point total; all shifts $p < 0.001$. 43

Table 7: Deployment measures by preset. API rows are priced from logged token use at July 2026 list prices; local rows are a total cost of ownership, measured electricity at a PUE of 1.54 plus a 1,600-dollar card over three years at half utilisation. The two are different kinds of number. Cost per accepted consultation divides by the panel accept rate in Table 4. 46

Table 8: Headline numbers with and without the two faulty cases. The 48-case column is primary throughout. 48

List of Figures

Figure 1: Overview of the study. A simulated clinical consultation and evaluation pipeline. (Top) Input eye case. Orchestrate clinical interactions between health worker agents and Alan overlayed on 07 LLMs.. (Middle) Generated interaction transcripts are aggregated into a central repository. (Bottom) Dialogue quality is evaluated by via parallel tracks: a panel of LLM judges and human clinical experts, both scoring responses against evaluation rubrics to yield comparative analytic metrics. 4

Figure 2: Annual cost of large language model deployment as a percentage of national health expenditure.....6

Figure 3: Example of a clinical reasoning trace in a conversational AI system. The health worker's observation is followed by the model's response and an explicit reasoning segment 10

Figure 4: WHO six ethical principles for the use of artificial intelligence for health 11

Figure 5: Schematic positioning map: evaluation realism against model accessibility.....13

Figure 6:Methodology overview – Part A-B - A)Build Stage: Preparation of frozen test cases B)Serve Stage: Dual deployment architecture routing requests either via external Frontier APIs (e.g., OpenRouter) or local inference on a High-Performance Computing (HPC).....15

Figure 7:Methodology overview- Part C - Workflow of the simulated consultation loop.....16

Figure 8: Methodology overview - Part D - Transcript validation, auditing, and anonymization pipeline.....17

Figure 9: Methodology overview - Part E - Automated evaluation pipeline using an LLM judge panel.....18

Figure 10: Methodology overview - Part F and G - F) Human validation anchoring and quantitative analysis pipeline G) Analyse: Downstream data synthesis integrating 1,400 dialogues, 8,400 AI judge panel scores, and 280 human ratings, as planned before the two case exclusions (the analysis set is 1,344, 8,064 and 268; see Section 3.1) 19

Figure 11: Alan turns per dialogue by preset. Box plots depict the distribution (median, interquartile range, and range) of the number of turns taken by Alan per dialogue 29

Figure 12: Mean words per Alan reply by preset. Box plots depict the distribution (median, interquartile range and range) of response lengths. The dashed horizontal line indicates the target scaffold length constraint (33 words). 30

Figure 13: Spearman rank correlation matrix of evaluation criteria ($n = 8{,}332$ $(8{,}064 + 268)$). Strong positive correlations are observed between C1 (Clinical Accuracy) and C2 (Urgency / Red-flag, $\rho = 0.70$), as well as between C4 (Clarity) and C5 (Scaffold Adherence, $\rho = 0.65$). C3 (LMIC Appropriateness) demonstrates weak correlation across all other criteria ($\rho \leq 0.23$). 31

Figure 14: Composite score against response latency. ($n = 1{,}344$ dialogues). Weighted composite score (0–100) plotted against total reply to latency in seconds on a logarithmic scale across evaluated models. A weak but statistically significant -ve correlation is observed overall (Spearman $\rho = -0.14$, $p = 5.7 \times 10 - 7$). 32

Figure 15: Model performance by criterion. Dot plots illustrate the average performance scores for each evaluated model across the five specific criteria, as well as the overall unweighted mean score. Values represent averages across thinking presets on and off. Concentric rings highlight the top-performing model within each evaluation panel, and the performance spread across models is detailed below each axis. 34

Figure 16: The three lowest- and three highest-scoring clinical scenarios, pooled over all 14 model presets and both worker personas (n = 1,344 dialogues). Blue markers show the mean weighted composite score (0 to 100), red markers the percentage of judgements passing the acceptance gate. Dotted lines mark the medians across all 48 cases 35

Figure 17: Model Capability Gradient (n = 1,344): Weighted composite scores (0–100) with 95% CIs across model presets. Highlights a top-tier plateau (80.7 - 84.2).....37

Figure 18: Paired thinking deltas per model with 95 percent CIs. ($n = 96$ per model). Bars depict the mean paired difference in weighted score (0–100 scale) for each model. Error bars represent 95% confidence intervals, with corresponding p-values shown above each bar.

Positive deltas indicate performance gains under extended thinking, while negative deltas indicate performance loss. 38

Figure 19: C1 Clinical Accuracy vs. C5 Scaffold Adherence (n = 1,344): Comparison of mean scores (1–5) across model presets ranked by overall capability. 39

Figure 20: Where models breakdown. (n = 8,064 judgements). Percentage share of judgements exhibiting hollow compliance (C5>= 4 and C1<=2 gold) versus format collapse (C5<= pink). 40

Figure 21: Health worker persona sensitivity across model presets: Performance gap between tailoring responses for a Community Health Worker (HW01) vs. a Clinical Officer (HW02) 41

Figure 22: Each confusion matrix illustrates the percentage and absolute counts of inter-evaluator decision overlap (accept vs. reject decisions) between model judges: (left) gpt-5.4-judge versus sonnet-5-judge, (middle) gpt-5.4-judge versus qwen3.6-27b and (right) qwen3.6-27b versus sonnet-5-judge. Colour intensity maps to the percentage of shared dialogues.....42

Figure 23: Agreement matrix between human evaluators (n = 134 shared dialogues). The percentage and absolute counts (n) of decision overlap (accept vs. reject) between Human 01 and Human 02. Colour intensity maps to the percentage of shared dialogues 43

Figure 24: Bland-Altman plot of the panel against the pooled physician mean on the 25 point total. The solid line is the mean difference, the dashed lines the 95 percent limits of agreement. 44

Figure 25: Break-even between owning a card and renting a frontier interface. The flat line is the local option; the sloped lines are cumulative API spend. 47

Figure 26: Judged quality against consultation time. Comparison of weighted composite quality scores (0–100 scale) against mean total Alan latency per consultation (seconds) across $n = 1{,}344$ simulated dialogues. 47

1. Introduction

1.1 Background and motivation

The global burden of vision impairment is large and inequitable. An estimated 1.1 billion people live with untreated vision impairment. Around ninety percent of it is avoidable and the burden sits overwhelmingly in low- and middle-income countries (LMIC) [1,2].

Eye specialists are scarce. At roughly 3.7 ophthalmologists per million people in LMICs against 76.2 per million in high-income countries [3]. The first contact for an eye problem is therefore a nurse, a clinical officer or a community health worker. Delegating care to such workers is deliberate policy. But knowledge from short trainings fades quickly. And single presenting clinical signs are too insensitive to support safe referral by checklist [4,5].

Multimodal foundation models, frequently referred in literature by the shorthand Large Language Models (LLMs) now demonstrate clinical knowledge and reasoning at or near specialist level on examination benchmarks, including eye care [6,7]. If that capability could be delivered reliably, safely and affordably to frontline health workers, it would support efforts to close the access gap.

1.2 Alan: an open-source clinical scaffold

Alan is an open-source clinical agent for eye, ear and skin care. It's built for frontline health workers in low-resource settings [8]. It is not itself a language model. It is a structured system prompt of roughly 15,000 words, which we call a scaffold. It can be laid over any model. The scaffold encodes a five-step clinical workflow running from history taking, through guided examination, to diagnosis, management and safety-netting. It enforces red-flag triage, steering the host to give urgent first aid guidance when there is a sight- or life-threatening condition. And it constrains every reply to a single short text, preferably 33 words and at most three questions. It responds in a language suitable for users with limited English [8].

The practical effect is that Alan is portable. Nothing in the scaffold is tied to a particular model. The same scaffold can be laid over a frontier online model reached through a commercial application programming interface (API), or over a small open-weight model running offline in the clinic itself.

That portability is what makes the questions in this dissertation answerable: the scaffold can be held still while everything underneath it changes.

1.3 The problem: model scale and the deployment gap

Informal testing suggests that Alan performs well on frontier models and weakens on small open-weight ones [9]. Small open-weight LLMs are more likely to omit mandatory triage steps, lose conversational context, violate the output format or misapply referral criteria. Frontier models are reached almost exclusively through proprietary and costly cloud interfaces. Frontier models depend on a live network connection. They charge for every consultation, so spending rises with patient volume. They send clinical conversations to servers outside the health facility or country. And it leaves the health system dependent on external pricing and availability for the life of the deployment [10,11].

Open-weight models potentially remove the above worries, after an initial hardware purchase. They run locally on anything from an institutional server to a consumer mobile phone. Clinical data never leaves the building. Health technologies should be designed for the contexts and capacities of low-resource settings rather than only for high-income ones [12]. What is not established is how much clinical quality that route gives up. Measuring that requires holding a scaffold fixed while the model beneath it changes. To our knowledge, no published work has done so.

1.4 Research questions

Three questions therefore remain open:

RQ1. How much of a fixed scaffold's clinical quality survive when it is moved from a frontier model to a small open-weight model that can run locally and where does it break down?

RQ2. Can a panel of LLM judges measure that difference reliably and does it agree with Human ophthalmologists?

RQ3. What would each LLM configuration cost to run at the frontline? and what quality is given up in exchange?

1.5 Significance of this study

The question sits squarely within health data science. Answering it required four disciplines at once, with clinical ophthalmology setting the cases and rubric, machine learning evaluation supplying the judge panel and its reliability statistics, high performance computing making the local models measurable, and health economics and regulation framing what the numbers are for.

This study offers a method as much as a result. Comparisons between clinical AI systems usually change everything at once, so it is not clear what caused what. Here only one thing changes. The scaffold instructions stay word for word the same and the model is swapped one after another. That turns a vague question into an answerable one. Not which model is best, but how much of the quality survives when the tool is moved onto a machine a low-resource clinic could buy. The design is portable. Any organisation with a prompt-based clinical workflow can run the same method against its own tool.

For practice, the study contributes a cost analysis that did not exist before. It shows what a single eye care consultation would cost and how long a health worker waits for each reply. The money side is indicative rather than exact, since it rests on price that each clinic would set differently.

For policy, this study reinforces the need for strict regulatory vigilance as generative AI enters healthcare. By demonstrating that identical prompt instructions yield different clinical behaviour when the model is swapped, this work highlights why regulatory frameworks must treat the prompt and model as an inseparable unit. The study underscores that safety authorizations evaluate a clinical scaffold and model architecture together as a single product.

1.6 Scope of the study

First, the clinical context covers a single specialty: eye care. Second, the data analysis evaluates text-based conversations only, excluding any multimodal communication. Finally, the financial assessment is restricted to an indicative cost estimate rather than a comprehensive health economic evaluation. An overview of the overall simulation and evaluation framework is displayed in ***Figure 1***.




Figure 1: Overview of the study. A simulated clinical consultation and evaluation pipeline. (Top) Input eye case. Orchestrate clinical interactions between health worker agents and Alan overlayed on 07 LLMs.. (Middle) Generated interaction transcripts are aggregated into a central repository. (Bottom) Dialogue quality is evaluated by via parallel tracks: a panel of LLM judges and human clinical experts, both scoring responses against evaluation rubrics to yield comparative analytic metrics.

2. Literature Review

2.1 Task shifting and the Arclight

Task shifting, delegating clinical tasks to less specialized workers, is the standard response to specialized health workforce scarcity. It's harder than it sounds. General health workers delivering primary eye care across Kenya, Malawi and Tanzania showed low eye-care knowledge, with training only weakly predicting performance [5]. Single presenting eye signs carry sensitivities of just 6 to 55 percent for referral-worthy disease, so no checklist can triage reliably [4]. Task-shifted workers in Sierra Leone found the model workable but the workers isolated [13]. Trained knowledge fades, single signs mislead and ongoing support is weak.

One of the evidenced responses so far has been hardware. The Arclight, a low-cost solar-powered ophthalmoscope designed for these settings. It let eye health workers in Malawi identify more clinical signs and make more correct diagnoses than a traditional instrument did [14–16]. In Tanzania, two facilitators trained 378 primary health workers through a cascade that screened around 36,000 children in six months [17]. And a follow-up study has tracked the long-term positive impact of the Arclight [18]. That programme solved a great deal of task shifting, that is about seeing the sign. The cognitive half remains open: deciding what the sign means, what to ask next and whether to refer. Knowledge that fades call for support at the point of care, rather than more courses and trainings. Signs that mislead argue for a conversational work-up rather than a flowchart. Alan (***Section 1.2***) has the potential to be a response to that.

2.2 The barriers to global digital health

Digital support for health workers was endorsed by the WHO, with the caution that the evidence was thin [19]. Reviews since then map the barriers, including internet access, cost, local language support and data control [20].

What separates LLMs from earlier digital tools is not just capability but also affordability. On 2025 prices, serving a 70 billion parameter model to a handful of concurrent users took roughly $300,000 a year, which in some LMICs would consume up to 15 percent of their national health spending per 100,000 people [21]. ***Figure 2*** shows how that scales with national income. Affordability is not a threshold but a slope. Where a health system sits determines which models are even candidates.




Figure 2: Annual cost of large language model deployment as a percentage of national health expenditure.

Source: Chen et al. (2025), Reproduced without modification under CC BY-NC 4.0.

How such costs should be judged is a separate literature and this study does not enter it. A health economic evaluation would express cost against a unit of health gained, counting staffing, training and downtime alongside hardware [22,23]. What ***Chapter 4 - Results*** report is narrower and comparison is reported as an indicative costing, not an economic evaluation.

2.3 RAG and commercial evidence tools as alternatives

One widely adopted strategy for enhancing medical applications of LLMs is Retrieval-Augmented Generation (RAG). RAG allows searching a library of papers and guidelines first, then writes its response with the sources attached [24]. Several products now work this way. OpenEvidence is one of the largest, with reported 40% of physicians in the U.S. utilizing the tool [25]. It is free to any doctor or medical student who has professional credentials in selected countries. It licenses content from journals such as NEJM, JAMA and Lancet [26]. UpToDate, another point-of-care, evidence-based tool, has anchored a similar AI layer onto its reference library [27].

A recent study compared OpenEvidence and UpToDate against three general purpose models (like ChatGPT and Google search). On exam questions, the best general-purpose model scored 97.4 percent, against 89.6 for OpenEvidence and 88.4 for UpToDate. Google's ordinary search summary scored as well as either of them. OpenEvidence came last of all six on clarity measures, which the authors read as a communication problem rather than a knowledge one [28]. All implying that fetching documents can make an answer worse, not better.

For LMIC setting trouble starts earlier. Every one of these tools needs a live internet connection and an account. The OpenEvidence account currently requires an American provider license number. The library it searches is built from American and European guidelines, so the answer arrives already committed to a level of care that may not exist in low-resource clinics. None of this makes RAG a bad idea. But it makes it a wrong move for an LMIC clinic with poor bandwidth. The alternative could be to carry the guidance in the prompt instead of fetching it.

2.4 The scaffold-model pair and instruction following

The assumption is that good clinical behaviour can be installed by text prompt alone. Studies support this premise. A structured clinical-reasoning prompt improved LLM's diagnostic accuracy on the Radiology Diagnosis [29]. And the CSEDB (Clinical Safety-Effectiveness Dual-Track Benchmark) team found structured prompting significantly improved both safety and effectiveness, reinforcing structured prompting as a reproducible enhancement method for clinical LLM reliability [30]. In other words, beneficial clinical behaviour is partly "promptable".

Products already in practice are built on that idea. Penda Health's AI Consult lays a decision support prompt over GPT-4o [31]. In Nairobi, across 39,849 clinic visits, it produced a 16 percent relative reduction in diagnostic errors and 13 percent in treatment errors [31]. The product runs on a host model and the host that carried Penda Health's result (GPT-4o) is already two generations old. A 16 percent reduction is therefore a property of a pair, not of a prompt alone.

This confounding is a gap in the literature. When a system is tested on benchmarks, the result comes back as a single score. That score covers two things at once: what the prompt contributes and what the model underneath can do on its own. There is no way to tell them apart. One review of thirteen AI coding assistants refused to compare their scores for exactly this reason. Each ran on a different model, and the choice of model shapes the result more than the design

of the instructions does [32]. Rombaut's recommended fix is a controlled experiment. Hold one part still and vary the other.

That fix has been tried outside medicine. A recent study on coding agents paired three scaffolds with five models and found that the scaffold alone shifted accuracy by as much as 28 points [33]. To our knowledge, nobody has yet tested this in clinical care. No study took a full clinical scaffold; kept it fixed and changed the model beneath it. That is the experiment this study runs.

Holding a behaviour across a conversation is hard. Laban et al. found that every model they tested performed, on average, 39 percent worse in multi-turn versions of tasks than in single-turn equivalents [34]. An interesting finding shows that models can restate the constraints and instructions they are currently violating. So the failure is more behavioural than of memory [35]. Another medical LLMs evaluation also exhibited severe vulnerabilities in conversational settings, with clinical accuracy dropping from 91.2% to as low as 13.5% for Claude Sonnet 4 model [36].

Alan, the scaffold in this study, is roughly 15,000 (~21,000 tokens). It has greeting, history-taking, red-flag checks, referral logic and safety-netting, that must hold across an entire consultation. All of it must survive a single conversation, from the first question to the last. The studies above suggest instructions this long start to slip as a conversation goes on. To our knowledge, no study yet measures whether the failure is uniform. A system can forget its instructions, forgetting what it was told, asking the same question twice and never settling on a diagnosis. Or it can follow every instruction, ask all the right questions and then recommend the wrong treatment. The distinction is clinically important. A model that follows instructions, while losing clinical judgement, would look safe to an adherence metric and be dangerous in clinical use.

The evaluation rubric in this study separates scaffold adherence from clinical accuracy. so that the two can be observed separately.

2.5 LLMs in eye care and their evaluation

Eye care adopted LLMs and evaluated them early. The pattern mirrors the wider field. In 2024, its reported LLMs show expert-level eye care knowledge in exam settings [7]. By 2026 an evaluation of ophthalmic emergencies compared GPT-5, Gemini 2.5 Pro and Claude Opus 4.1 on 40 emergency vignettes and found diagnostic accuracies above 80 percent with no statistically

significant differences between the models [37]. A study of deep-reasoning models on staged, workflow-mimicking information levels found overall triage accuracy of 85.3 percent [38]. Another 2026 study tested five frontier models on 3,831 real referral vignettes written in Portuguese and reported 68.7 percent baseline triage accuracy, improving with labelled examples [39].

The literature has three key limitations. First, evaluations conducted are single-turn or staged, giving LLMs all the facts upfront rather than forcing it to ask questions and gather a patient's history. Second, almost all studies focus on top-tier, proprietary models that rural clinics cannot run. Third, these studies ignore real-world constraints, scoring recommendations against high hospital standards without considering what is actually available in low-resource eye clinics. And if frontier models are statistically indistinguishable at the top of the range, then the useful question is not which model wins. It is how far down the scale good consultation survives

2.6 LLM-as-a-judge, rubrics and the reasoning

Scoring thousands of clinical consultations by human experts is not feasible. The field therefore increasingly uses other LLMs as judges. The approach can have alignment with human evaluation on open-ended tasks [40,41]. A judge LLM is good enough when it agrees with humans about as often as humans agree with each other [42].

Limitations of using LLMs as a judge is equally well documented. They tend to reward long answers, documented as “verbosity bias” [43,44]. They also have “position bias”, as bias caused by respondents' tendency to opt for alternatives in certain positions (e.g., the first thing mentioned) regardless of their content [45,46]. Another one is “self-preference bias”. It’s demonstrated that LLM evaluators tend to score their own generated outputs higher than outputs from other models or humans, even when quality is equal [47]

Healthcare has taken the method up. One analysis showed LLM-human judgement alignment (Cohen’s kappa of 0.59 to 0.93) and alignment strongest where the task was tightly scoped and the rubric explicit [48]. Two 2026 papers complicate that. A frontier judge reached the physician agreement ceiling while showing no clinical caution at all, scoring confidently on the very items where physicians held back [49]. And LLM judges reached the same verdicts as humans by different reasoning, so matching scores do not imply matching reasons [50]. LLM-as-a-judge is a measurement instrument, and instruments need calibrating before their readings are trusted.

2.7 Chain-of-thought reasoning as diagnostic evidence

Researchers have moved beyond simple response generation by introducing the concept of "thought", a sequence of tokens (words, phrases or punctuations that an LLM reads, processes and generates) representing intermediate steps in the reasoning process. This innovative paradigm enables LLMs' to mimic complex human reasoning processes [51,52]. An example of what an LLMs chain of thought is shown in ***Figure 3.***




Figure 3: Example of a clinical reasoning trace in a conversational AI system. The health worker's observation is followed by the model's response and an explicit reasoning segment

There are certain concerns on reasoning models. One is faithfulness of the models. Reasoning models verbalise the hints they actually used only about 25 to 39 percent of the time [53]. Furthermore, verbalized reasoning can give an incorrect picture of how models land at conclusions [54]. These "thinking" traces are therefore evidence of behaviour, not much use as proof of how the model works. Therefore, the exploratory analysis in this study reads them accordingly.

For frontline deployment, the question is sharpened by cost. "Thinking on" is paid for in seconds of latency and in energy. Both are scarce in the settings this study targets. LLM response

that arrives two minutes late may be no gain at all in a busy clinic. The thinking toggle is therefore a deployment decision with price.

What literature lacks here is a controlled study. The same cases, the same scaffold, thinking toggled on and off, across a ladder of models. This study design provides one.

2.8 Regulation and governance of clinical LLMs

The governance frame for digital health is set by the WHO. Its 2021 guidance on the ethics of AI for health set out six principles [12]. Its 2024 guidance on LLMs applied those principles and issued more than forty recommendations, calling explicitly for LMIC participation in design and governance [55]. ***Figure 4*** sets out the six principles.



Figure 4: WHO six ethical principles for the use of artificial intelligence for health

Source: World Health Organization (2024), Reproduced without modification under CC BY-NC-SA 3.0 IGO.

Challenging questions were directed at the WHO frame. Can AI serve universal health coverage in LMICs at all? given who builds these systems and where the data goes [56].

Offline, open-weight deployment answers part of it. A consultation never leaves the device, and no third party holds the record, which speaks to *autonomy*; connectivity requirements and per-use fees disappear, which speaks to *inclusiveness and equity*; and a pinned version behaves the same next year as this year, so a failure can be traced rather than guessed at.

The limits are equally real. Open weights are not an open explanation: without training code and data the work is harder to check than the word "open" suggests [57,58]. None of it reaches the health worker, who still cannot see why the answer came out as it did. And current explanation methods give partial accounts rather than safety guarantees [59]. Going offline is not free either, with close to a billion people served by facilities that have no reliable electricity [60].

This study highlights a challenge for medical AI safety. Regulators know how to approve software that changes after release under a pre-specified plan [61,62]. But with AI, the product is often split in two: one group writes the clinical rules (the scaffold, like Alan), the another updates the underlying AI model. When a life-threatening medical error occurs, which half of the product failed?

2.9 The gap this study occupies

Existing research often misses how real-world eye care works. Studies give AI models all the patient facts upfront, but real healthcare workers must extract those facts themselves. While conversational AI tests exist, they use unguided models. Almost every study evaluates models using high-resource hospital standards. Limited work has tested a single structured clinical scaffold across a range of model sizes, with thinking on and off, to see where it breaks, how well it performs or what it costs. Finally, it remains unsettled if LLM judges can accurately evaluate these tools. ***Figure 5*** maps this study within other literature.




Figure 5: Schematic positioning map: evaluation realism against model accessibility

Conceptual positioning of this study within the clinical LLM evaluation literature. The vertical axis runs from static exam formats to multi-turn dialogue resembling real clinic workflow. The horizontal axis runs from proprietary cloud-only models to open-weight models that can be run locally. Positions are indicative and show relative ordering only. Neither axis is a measured scale, and no quantitative comparison between studies is intended. Author's own figure, based on the studies reviewed in this chapter

3. Methodology

3.1 Methodology overview

This study is a simulation experiment. Eye cases were played out as full consultations between a simulated frontline health worker and Alan. The scaffold was hosted on seven models in turn: two frontier systems, GPT-5.4 [63] and Claude Sonnet 5 [64], top commercial releases available at the time [65], and five open-weight models in ***Table 3*** . Each ran twice, with reasoning ("thinking") ON and OFF giving fourteen presets. Each preset met both worker personas on all cases.

Every dialogue was scored on a five-criterion rubric by three judge models, each playing two expert personas: six independent judgements per dialogue. Two ophthalmologists independently scored the same 10 percent subsample to anchor the panel.

Two cases were then removed. Their reference diagnosis and management had been transposed with each other during data entry. The human ophthalmologists found this during validation (***Section 3.4***). The analysis set is therefore 48 cases, 1,344 dialogues, 8,064 panel judgements and 134 physician-scored dialogues.

Each judgement was reduced to a weighted composite score and an acceptance decision (***Section 3.3***). The panel was treated as an instrument against itself and against human ophthalmologists. Energy and latency cost were measured. ***Figure 6*** to ***Figure 10*** summarise the technical pipeline.

All data, code, dialogues, scores and full pipeline can be found in public ***GitHub repository***.




Figure 6:Methodology overview – Part A-B - A)Build Stage: Preparation of frozen test cases B)Serve Stage: Dual deployment architecture routing requests either via external Frontier APIs (e.g., OpenRouter) or local inference on a High-Performance Computing (HPC)




Figure 7:Methodology overview- Part C - Workflow of the simulated consultation loop




Figure 8: Methodology overview - Part D - Transcript validation, auditing, and anonymization pipeline




Figure 9: Methodology overview - Part E - Automated evaluation pipeline using an LLM judge panel




Figure 10: Methodology overview - Part F and G - F) Human validation anchoring and quantitative analysis pipeline G) Analyse: Downstream data synthesis integrating 1,400 dialogues, 8,400 AI judge panel scores, and 280 human ratings, as planned before the two case exclusions (the analysis set is 1,344, 8,064 and 268; see Section 3.1)

3.2 Simulation as the evaluation method

One practical way to test a scaffold like Alan is to simulate consultation workflow. Our goal was to quantify the fall in quality across models. In a real clinic, a study large enough to separate seven models would run for years.

In-silico testing is not new. AMIE was tested in simulated consultations against primary care physicians, with trained patient actors on the other side [66]. CRAFT-MD simulated multi-turn conversation and found that models which ace exam questions fall apart once they have to take the history themselves [67]. That second result is the reason this design is a conversation and not a questionnaire or MCQ.

The limits of simulation studies are known. A simulated health worker never gets tired and never gets interrupted. We took that trade knowingly. Alan is not a validated tool. And a field study would put it between a health worker and a real patient before anyone knows its failures and limitations. This is not to say the field trial can be skipped.

3.3 Evaluation rubric

Every consultation is scored on five criteria, each anchored on a 1 to 5 scale. ***Table 1*** gives each criterion, its weight and the reason it was chosen. The full rubric, with worked anchors for every score point, is in ***Appendix II: Evaluation Rubric***.

Table 1: The evaluation rubric, its weights and the reason each criterion was included.

Criterion/Weight	Question it answers	Rationale
C1: Clinical accuracy, 30%	Was a clinically defensible diagnosis, differential ranking and management reached?	Diagnostic accuracy is the foundational dimension of clinical AI evaluation [68,69].
C2: Urgency Calibration /red flag detection, 30%	Were all red flags identified and escalated at the right urgency?	In ophthalmic emergency evaluation, failure to recognise red flags can lead to devastating, and sometimes fatal, consequences for the patient [37]
C3: LMIC appropriateness, 20%	Is recommendation feasible in a basic rural clinic?	Operational feasibility, rather than accuracy alone, determines whether clinical AI can be adopted in a low-resource setting [12,21]. ***Appendix VI: LMIC medication and instruments*** reference adopted from WHO package of eyecare interventions [106]

Criterion/Weight	Question it answers	Rationale
C4: Communication clarity, 10%	Could a non-specialist understand and act on this guidance?	HealthBench lists communication as a core evaluation axis [68]. CLEVER found conciseness separating from factuality under blind clinician review [69].
C5: Scaffold adherence, 10%	Did the consultation follow the scaffold's clinical sequence?	A diagnosis reached by skipping or reordering steps is difficult to audit or safely follow.

Every rater gives five scores, one per criterion, each from 1 to 5. Two numbers are then built from them, the same way for every rater.

The first is the weighted composite. Each score is multiplied by its criterion's weight and the five results are added. That gives a number between 1 and 5, which is put on a 0 to 100 scale by subtracting 1, dividing by 4, and multiplying by 100. The composite is worked out one judgement at a time. Criteria are never averaged across raters first. So, a figure quoted for a preset is the average of that preset's individual composites.

The second is the acceptance gate. A straight pass or fail. A rater accepts a consultation when clinical accuracy (C1) and urgency calibration (C2) are both 4 or more, and LMIC appropriateness, communication clarity and scaffold adherence (C3 to C5) are each 3 or more. All five conditions must hold together. A consultation that is clear and well-ordered but reaches the wrong diagnosis therefore fails, however high those other scores are. Each dialogue is judged six times, and the accept rate is the share of all its judgements that passed. It is not a majority verdict, so disagreement between judges is kept rather than resolved away.

Accuracy and urgency calibration carry 30 percent each, because either can lead directly to avoidable blindness. LMIC appropriateness carries 20 percent, because advice that cannot be delivered is not advice. Communication clarity and scaffold adherence carry 10 percent each. No elicitation study supports that split, and ***Section 4.10*** reports what moving any weight by ten points does to the ranking.

3.4 Case bank

Public eye case sets exist and the method would work with any of them [67,70,71], but none restricts its findings and management options to what a low-resource clinic can do.

A case bank was assembled for this study by eye care specialists with clinical and teaching experience in low-resource practice. Every record was written against what a rural health post can see and do. The 50 cases fall into five groups: loss of vision (13), acute trauma (11), red eye (10), lumps and bumps (8) and paediatrics (8). Each holds an opening line, age, sex, symptoms, duration, history, five examination fields and a reference diagnosis and management plan see ***Table 2*** for example case presentation.

Table 2: Example record from the case set, case 01.

Diagnosis	Cataract
First Line for Alan	Old man, outside work. Vision poor. Shadows in pupil.
Management	surgery
Age	65
Sex	m
Symptoms	glare
Duration	years
Drugs, history, job etc	outdoor jobs
Visual Acuity	6/60, 6/36
Pupils	OK
Front of Eye & Lids	OK
Fundal Reflex	hazy, spokes
Back of Eye	good

The worker agent holds those facts but reveals a finding only when Alan asks. The reference ground truth diagnosis and management is shown only to the judges. Judges mark against a key rather than solving the case themselves. That lowers the reasoning burden on the judges.

Two cases did not survive contact with the physicians. Cases 48 and 49 had their reference answers transposed with one another, so neither could be marked fairly. Therefore, a 48-case set is primary data throughout the study. Worth noting, the two human ophthalmologists found this error when neither the judge panel nor any automated check did. ***Table 8*** reports a sensitivity check if all 50 cases were consumed .

3.5 The scaffold

Alan is an open-source structured system prompt of roughly 15,000 words (21,000 tokens) [8]. One compiled version was frozen for the whole study. A snapshot of what Alan looks like is in ***Appendix V: Snapshot of Alan, the scaffold.***

3.6 The simulated health worker

One model, Claude Sonnet 5 [64], roleplays health worker in two personas. HW01 (health worker 01) is a community health extension worker at a rural health post, with very limited English and basic clinical training; HW02 is a district clinical officer with confident clinical English. Each prompt fixes setting, equipment, vocabulary, style and behavioural rules. The two are a factor rather than a robustness check. ***Section 4.7*** reports the effect. Both prompts are in ***Appendix III: Health Worker Persona System Prompts***.

Role-based prompting is the standard way to build such an agent [72]. One capable model, Claude Sonnet 5, was used throughout. If the worker changed with the host, worker LLM quality would be tangled up with host quality. Similar to pilot runs we conducted, less capable models drift out of persona and break their rules [73].

That choice has a limitation as well. Claude Sonnet 5 plays the worker in all 1,344 dialogues while also serving as an Alan host and as one of the three judges. The worker-to-host overlap means the comparison between Claude Sonnet 5 as a host and the others is not clean [47].

Worker temperature was set to 0.7, following common practice for this kind of roleplay [74,75]. The worker is not scored and takes no clinical decision.

3.7 The LLM judge panel

A single judge imports one model's biases, while a panel of diverse judges outperforms one large judge and averages out intra-model bias [76]. The panel crosses three judge models from three families with two expert personas, so every dialogue receives six independent judgements. E1 (expert 1) is a consultant ophthalmologist with rural programme experience. E2 is a health educator who trains community health workers. Both use the same rubric and score all five criteria. Full expert persona prompts can be found in ***Appendix IV: Expert Judge Persona System Prompts.***

Judge temperature is set at 0.1, because a judge is a measuring instrument and low temperature gives the most consistent judgements. Though determinism alone does not guarantee validity [77]. That is why the human anchor was added.

3.8 Human expert validation

Two ophthalmologists independently scored the same subsample. Both are practising specialists with extensive clinical experience in low-resource ophthalmology and global eye care. The standard the LLM panel is calibrated against is therefore set by clinicians who have worked in the kind of setting the scaffold was written for. Both scored every dialogue, so human-to-human and human-to-LLM-panel agreement rest on identical material.

Both were given the rubric, its anchors and a worked walkthrough of the criteria and the gate before scoring. They then scored independently and remotely, across several sittings, seeing neither the other's scores nor any model identity. They read the same anonymised transcripts the LLM judges read (***Section 3.10***). Presentation order was randomised. A physician figure quoted against the LLM panel is the mean of the two raters' scores.

The sample is a proportionate stratified random draw [78] with the model-by-persona cell as the stratum.

Three kinds of answer need three measures. For the 1 to 5 scores and their total, the primary statistic is ICC(2,1) [79], which asks whether two raters wrote the same number rather than merely ranked cases alike. Quadratic-weighted kappa beside it [80] and Krippendorff's ordinal alpha covering the ragged eight-rater matrix [81].

For the accept decision we report raw agreement, Cohen's kappa and Gwet's AC1, since kappa reads badly when almost every case falls on one side [82,83]. Bland-Altman limits can show steady bias between raters [84].

A panel is good enough when its agreement with humans is comparable to the agreement humans reach with each other [42]. The full-overlap design makes that possible, because the human-to-human figure on the same 134 dialogues is the ceiling. ***Section 4.9*** reports both sides.

3.9 Dialogue orchestration and environment

Each dialogue runs as a loop. The worker opens with the case's first line, Alan responds, and the worker answers. A conversation ends in three scenarios. One, when Alan produces its closing "Step 5 reply", detected by its fixed closing phrases. Second, at a hard cap of sixteen Alan turns, since pilot runs showed longer conversations were looping. ***Section 4.1*** reports how often each preset ended at the cap. And third, when there is no reply from Alan, for technical reasons. In such case, a dialogue will be reinitiated.

Every transcript records the conversation, the model's reasoning (kept separate and never shown to the worker or judges), latency and tokens.

Alan's temperature is fixed at 0.3 for every host. Across nine models and a range of benchmarks, temperatures from 0.0 to 1.0 produce no significant difference [85]. The generation budget is 8,192 tokens per turn, sized so reasoning models can finish thinking and still answer. Context window is 64K against an observed peak of roughly 40K.

All local inference ran on the gpu.A100 partition of the University of St Andrews Hypatia cluster [86], scheduled through Slurm 23.02.6 [87]. Each node carries four NVIDIA A100-SXM4-80GB GPUs, two AMD EPYC 7713 processors and roughly 1 TB of RAM, on Rocky Linux 9.4 with driver 555.42.06 and CUDA 12.5 [88]. Every dialogue job used a single A100.

Models were served as local OpenAI-compatible endpoints by three backends chosen by format: SGLang for bf16, llama.cpp for the q4_0 GGUF files, and a containerised transformers shim under Apptainer 1.1.0 where a model's architecture was newer than the cluster driver. Frontier hosts, the worker and the two API judges, were reached through OpenRouter [89]. ***Table 3*** lists every model.

Table 3: Every model used, with source, serving backend and settings

Role	Model	Source (accessed July 2026)	Serving / precision	Sampling and settings
Alan host	GPT-5.4	OpenAI[63]; OpenRouter: openai/gpt-5.4	API	T 0.3; 8,192 tok/turn; think on = effort high; off = minimal tier*
Alan host	Claude Sonnet 5	Anthropic[64]; OpenRouter: anthropic/claude-sonnet-5	API	T 0.3; 8,192 tok/turn; think on/off
Alan host	Gemma 4 31B	Google[90]; HF: google/gemma-4-31B-it-qat-q4_0-gguf	llama.cpp; QAT q4_0 (~18 GB)	T 0.3; 8,192 tok; 64K ctx; repeat penalty 1.1; top-k 64; thinking flag on/off
Alan host	Gemma 4 26B-A4B (MoE, 3.8B active)	Google [90]; HF: google/gemma-4-26B-A4B-it-qat-q4_0-gguf	llama.cpp; QAT q4_0 (~15 GB)	Same as Gemma 4 31B
Alan host	Gemma 4 12B	Google [90]; HF: google/gemma-4-12B-it-qat-q4_0-gguf	llama.cpp; QAT q4_0 (~7 GB)	Same as Gemma 4 31B
Alan host	Gemma 4 E4B	Google [90]; HF: google/gemma-4-E4B-it-qat-q4_0-gguf	llama.cpp; QAT q4_0	Same as Gemma 4 31B
Alan host	Qwen3-8B	Qwen[91]; HF: Qwen/Qwen3-8B	SGLang; bf16 (~16 GB)	T 0.3; 8,192 tok; 64K ctx; native thinking on/off
Health worker	Claude Sonnet 5	Anthropic[64]; OpenRouter: anthropic/claude-sonnet-5	API	T 0.7; 300 tok/reply; reasoning off
Judge	GPT-5.4	OpenAI[63]; OpenRouter: openai/gpt-5.4	API	T 0.1; reasoning minimal*; guided JSON schema
Judge	Claude Sonnet 5	Anthropic[64]; OpenRouter: anthropic/claude-sonnet-5	API	T 0.1; guided JSON schema; rubric prompt-cached; reasoning off
Judge	Qwen3.6-27B	Qwen[92]; HF: Qwen/Qwen3.6-27B	hf-shim (Transformers); bf16 (~54 GB)	T 0.1; 32K ctx; guided JSON; thinking off

HF = Hugging Face repository (Hugging Face, 2026), T = temperature, tok = tokens, ctx = context window
**GPT-5.4 cannot turn reasoning off entirely, so off means minimal.*

3.10 Blinding and bias controls

Judges never see who they are judging. Each transcript is anonymised first. Participants are relabelled "health worker" and "advisor" and model names, reasoning traces, timings and tooling metadata are stripped. An automated leak scanner then blocks any transcript that still identifies a model or reveals that a participant is an AI. Our run caught four incidents where the host named itself mid-dialogue [47].

3.11 Statistical Analysis

Analysis was done in a single Jupyter notebook in Python 3.13.14 [93] . The primary outcomes are the weighted composite on 0 to 100 and the acceptance rate (***Section 3.3***).

Uncertainty comes from a nonparametric bootstrap of 2,000 resamples. We resampled cases rather than judgements [94], because cases are the sampling unit and the six judgements of one dialogue are not independent. All intervals are 95 percent percentile intervals under one recorded seed.

Thinking-On vs Off effects are estimated within model as paired differences on the same case and persona, using Wilcoxon signed-rank tests [95]. The worker persona is treated the same way, as a second experimental factor paired across the same 48 cases and hosts. Associations such as score against reply length use Spearman correlations [96]. Ranking stability is stress-tested by recomputing the leaderboard under alternative gates and weights, reporting Kendall's tau, and by leave-one-judge-out re-analysis [97].

Seven paired thinking comparisons are made, one per model and they are the only family of formal tests; everything else is an estimate with an interval. A Bonferroni correction across those seven moves the threshold from 0.05 to 0.0071, and ***Section 4.5*** reports the raw p values.

Differences between presets are reported as estimates with intervals rather than as tests against a threshold, because no threshold exists to test against. Calling one preset close enough to another requires a statement about how much clinical quality a health system should trade for offline operation and lower cost, which is a policy judgement rather than a statistical one. ***Section 4.4*** and ***Table 5*** reports each shortfall with a case-clustered interval, then counts how many presets fall within a range of candidate thresholds, the leader included.

3.12 Energy footprint and running cost estimation

Throughout the local runs each card was logged with nvidia-smi every five seconds [98], and for every consultation the power curve was integrated across its start and end time. Idle draw was separated by the utilisation reading. The two proprietary models cannot be measured, so they are priced from logged token use.

Turning energy into money took three assumptions. Electricity is 26.11 pence per kWh, the Ofgem cap for July to September 2026 [99], converted at 1.3385 dollars to the pound. Facility overhead is a power usage effectiveness (PUE) of 1.54, the global average in the Uptime Institute's 2025 survey [100]. A clinic workstation would sit below that, so the local figure is conservative. Hardware is a 24 GB card at an indicative 1,600 dollars, spread across three years at half utilisation.

The two figures are not the same kind of number. The local figure is a cost of ownership: measured electricity plus amortised hardware for a machine the clinic owns. The frontier figure is a vendor's list price, which already contains that vendor's hardware, energy, support and margin. The local figure also omits networking, cooling, spares, technician time and downtime after a power cut. Any ratio indicates an order of magnitude rather than a like-for-like account.

3.13 Reproducibility and data availability

Every transcript carries model identifiers, seeds and timestamps. The case bank, personas, rubric, orchestration code, power logs and analysis notebook are versioned. The sampling script reproduces the human subsample exactly from its recorded seed. All available in public ***GitHub repository***.

3.14 Ethics

No patients and no patient data are involved, and all clinical content is synthetic. The only human participants are the two ophthalmologist raters, who scored anonymised dialogues and took no clinical decisions. Ethical approval was obtained through the School's ethics process and is reproduced in ***Appendix I: Ethical Approval***. This study establishes what the models do in simulation. It does not establish how the tool performs in the hands of real health workers.

4. Results

4.1 Dialogues

The run produced 1,344 dialogues: 48 cases by 14 presets by 2 worker personas. The median consultation took 12 turns, six from Alan and six from the worker. The frontier models and Gemma 4 31B held their replies close to the scaffold's 33-word target, averaging 32 to 35; Gemma 4 26B-A4B with thinking on drifted to 72. ***Figure 11*** and ***Figure 12*** give the distributions.




Figure 11: Alan turns per dialogue by preset. Box plots depict the distribution (median, interquartile range, and range) of the number of turns taken by Alan per dialogue

How a consultation ended separates the models more sharply than how long it ran. Qwen3-8B closed dialogue without naming a diagnosis in about a third of its consultations; every other preset named one at least 81 percent of the time. At the floor, the scaffold is not producing a poor consultation, it is failing to produce one at all. Details can be found in ***Appendix VII: Dialogue completion by preset.***




Figure 12: Mean words per Alan reply by preset. Box plots depict the distribution (median, interquartile range and range) of response lengths. The dashed horizontal line indicates the target scaffold length constraint (33 words).

4.2 Score data

The three LLM judges returned 8,064 judgements and the two physicians added 268 ratings. Depending on the criterion, 27 to 49 percent of judgements sit at the ceiling of 5 and floor scores are rare. C2 (urgency) and C5 (adherence) carry the highest means at 4.05, C1 (accuracy) the lowest at 3.83. The five criteria correlate without collapsing into one another. C3 (LMIC appropriateness) is the outlier: it correlates 0.10 with accuracy and 0.10 with adherence, against 0.70 between accuracy and urgency. ***Figure 13*** shows the correlation matrix.




Figure 13: Spearman rank correlation matrix of evaluation criteria ($n = 8{,}332\ (8{,}064 + 268)$)***.*** *Strong positive correlations are observed between C1 (Clinical Accuracy) and C2 (Urgency / Red-flag,* $\rho = 0.70$*), as well as between C4 (Clarity) and C5 (Scaffold Adherence,* $\rho = 0.65$*). C3 (LMIC Appropriateness) demonstrates weak correlation across all other criteria* ($\rho \leq 0.23$*).*

At dialogue level, quality falls slightly as consultations get longer (Spearman rho = -0.36, p < 0.001) and marginally as they get slower (rho = -0.14, p < 0.001; n = 1,344). ***Figure 14*** plots the latency relationship.




Figure 14: Composite score against response latency. ($n = 1{,}344$ dialogues). Weighted composite score (0–100) plotted against total reply to latency in seconds on a logarithmic scale across evaluated models. A weak but statistically significant -ve correlation is observed overall (Spearman $\rho = -0.14$, $p = 5.7 \times 10^{-7}$).

4.3 Leaderboard

GPT-5.4 with thinking on leads with a mean weighted composite of 84.2 (95 percent CI 80.7 to 87.3), and 71.9 percent of its judgements cleared the acceptable gate defined in ***Section 3.3***. Qwen3-8B holds the floor at 51.4. ***Table 4*** lists all scores.

Scores are from the LLM panel only; the ophthalmologists' ratings are held back for ***Section 4.9*** and reported as a sensitivity check in ***Table 8.***

The AI judge panel marks systematically harder than the two ophthalmologists. On the same 134 dialogues, the physicians accepted 79.8 percent and the LLM panel 54.1 (***Section 4.9***). Every accept percentage below is therefore a lower bound on what a clinician would accept and an instrument reading rather than a clinical verdict.

The LLM scores order the models reliably.

Table 4: Leaderboard of all fourteen presets, with case-clustered bootstrap 95 percent intervals on the weighted composite. Accept % is the share of each preset's 576 judgements that cleared the gate.

Preset	Weighted 0-100	95% CI	Accept %
GPT-5.4[on]	**84.2**	80.7 to 87.3	**71.9**
Claude Sonnet 5[off]	82.8	79.0 to 86.2	70.8
GPT-5.4[off]	82.2	77.6 to 86.3	71.0
Claude Sonnet 5[on]	82.1	78.4 to 85.8	69.3
Gemma 4 31B[on]	**81.8**	78.3 to 84.7	**67.9**
Gemma 4 31B[off]	80.7	77.9 to 83.5	62.5
Gemma 4 26B-A4B[off]	77.0	72.2 to 81.5	61.3
Gemma 4 12B[on]	77.0	72.8 to 80.7	57.8
Gemma 4 26B-A4B[on]	74.9	69.9 to 79.6	56.1
Gemma 4 12B[off]	69.6	64.3 to 75.0	44.1
Gemma 4 E4B[on]	65.1	59.9 to 70.1	43.9
Gemma 4 E4B[off]	59.1	54.0 to 64.7	31.8
Qwen3-8B[off]	53.4	47.3 to 59.2	26.2
Qwen3-8B[on]	**51.4**	45.7 to 56.9	**24.0**

The leading preset scored 4.59 on urgency and 4.64 on adherence but 3.87 on LMIC appropriateness. Across all fourteen, C3 spans only 3.57 to 4.16, the narrowest range of the five. ***Figure 15*** shows the pattern.




Figure 15: Model performance by criterion. Dot plots illustrate the average performance scores for each evaluated model across the five specific criteria, as well as the overall unweighted mean score. Values represent averages across thinking presets on and off. Concentric rings highlight the top-performing model within each evaluation panel, and the performance spread across models is detailed below each axis.

Case difficulty spreads widely. Onchocerciasis (Case06) is hardest with fewer than 2 percent of judgements accepting it, then trachoma (Case12) at 49.7. At the other end blepharitis reaches 91.2 and presbyopia 90.9, both accepted about nine times in ten. ***Figure 16*** gives the atlas. ***Section 5.4*** takes up what the two hardest cases have in common.




Figure 16: The three lowest- and three highest-scoring clinical scenarios, pooled over all 14 model presets and both worker personas (n = 1,344 dialogues). Blue markers show the mean weighted composite score (0 to 100), red markers the percentage of judgements passing the acceptance gate. Dotted lines mark the medians across all 48 cases

4.4 Frontier to local gap

The best frontier preset beats the best local one by 2.5 weighted points (95 percent CI 0.3 to 4.8***). Figure 17*** displays the full range . On the acceptance gate the same comparison gives 4.0 percentage points (CI -2.8 to 10.8), and that interval contains zero. Both ends of this comparison are the top scoring preset in their class, chosen from the same data used to estimate the gap. The local maximum is drawn from ten presets and the frontier maximum from four, so the local figure carries more of that selection advantage, and 2.5 points is best read as a floor on what a fresh case set would show rather than an unbiased estimate.

Every preset's shortfall against the leader is displayed in ***Table 5***, paired case by case. Three configurations cannot be separated from GPT-5.4 with thinking on at all: Claude Sonnet 5 with thinking off (1.5 points, CI -2.0 to 4.8), GPT-5.4 with thinking off (2.0, -0.3 to 4.3) and Claude Sonnet 5 with thinking on (2.1, -1.0 to 5.1). Gemma 4 31B with thinking on is the closest local configuration and the first whose interval clears zero, at 2.5 points (0.3 to 4.8). On the acceptance gate, the intervals are wider and four presets besides the leader contain zero.

Table 5: Shortfall against the leading preset (GPT-5.4 with thinking on), paired case by case, with case-clustered bootstrap intervals.

Preset	Composite shortfall	95% CI	Accept-gate shortfall (pp)	95% CI
GPT-5.4 [on]	leader		leader	
Claude Sonnet 5 [off]	1.5	-2.0 to 4.8	1.0	-7.3 to 9.0
GPT-5.4 [off]	2.0	-0.3 to 4.3	0.9	-5.6 to 7.3
Claude Sonnet 5 [on]	2.1	-1.0 to 5.1	2.6	-6.4 to 11.6
Gemma 4 31B [on]	**2.5**	**0.3 to 4.8**	**4.0**	**-2.8 to 10.8**
Gemma 4 31B [off]	3.5	0.5 to 6.7	9.4	1.6 to 17.4
Gemma 4 26B-A4B [off]	7.2	3.2 to 11.2	10.6	1.4 to 20.0
Gemma 4 12B [on]	7.2	3.0 to 11.3	14.1	3.1 to 24.7
Gemma 4 26B-A4B [on]	9.3	5.6 to 13.4	15.8	7.1 to 24.5
Gemma 4 12B [off]	14.6	9.4 to 19.8	27.8	16.5 to 38.9
Gemma 4 E4B [on]	19.1	14.4 to 23.9	28.0	17.4 to 38.5
Gemma 4 E4B [off]	25.1	20.3 to 29.6	40.1	28.3 to 51.4
Qwen3-8B [off]	30.9	24.7 to 37.2	45.7	31.4 to 58.9
Qwen3-8B [on]	32.8	27.1 to 38.6	47.9	35.9 to 59.7




Figure 17: Model Capability Gradient (n = 1,344): Weighted composite scores (0–100) with 95% CIs across model presets. Highlights a top-tier plateau (80.7 - 84.2)

4.5 Thinking on vs thinking off

Extended thinking pays off for two models only. Gemma 4 12B gains 7.4 points (CI 2.9 to 11.9, Wilcoxon p = 0.003) and Gemma 4 E4B gains 6.1 (CI 2.6 to 9.7, p = 0.004). Both survive a Bonferroni correction across the seven paired tests, though not by much: the corrected threshold is 0.0071. No other model moves beyond noise and three trend downward. ***Appendix VIII: Paired effect of thinking on*** gives every model's paired difference and ***Figure 18*** plots them.




Figure 18: Paired thinking deltas per model with 95 percent CIs. ($n = 96$ per model). Bars depict the mean paired difference in weighted score (0–100 scale) for each model. Error bars represent 95% confidence intervals, with corresponding p-values shown above each bar. Positive deltas indicate performance gains under extended thinking, while negative deltas indicate performance loss.

4.6 Adherence against accuracy

C1 asks if diagnosis and management response was right and C5 whether the scaffold's sequence was followed. Reporting them side by side shows which of the two survives further down the ladder.

Adherence outscores accuracy in thirteen of the fourteen presets as depicted in ***Figure 19.*** The single exception being Qwen3-8B with thinking on, where adherence falls 0.59 below accuracy.




Figure 19: C1 Clinical Accuracy vs. C5 Scaffold Adherence (n = 1,344): Comparison of mean scores (1–5) across model presets ranked by overall capability.

Hollow compliance, judgements in which the scaffold was followed well but the advice was wrong, peaks at 25.3 percent for Gemma 4 12B with thinking off.

Format collapse, where the scaffold broke down altogether, behaves in the opposite way. All presets stay under 3 percent for every preset scoring above 75. ***Appendix IX: Clinical accuracy against scaffold adherence*** and ***Figure 20*** give both counts for every preset.




Figure 20: Where models breakdown. (n = 8,064 judgements). Percentage share of judgements exhibiting hollow compliance (C5>= 4 and C1<=2 gold) versus format collapse (C5<= pink).

4.7 The health worker personas

Consultations with health worker 01 (limited resource and limited English) score 71.8 against health worker 02 (clinical officer, better English and better setup) 74.1; a difference of 2.3 points. Details can be found in ***Appendix X: Weighted composite by health worker persona.***

The persona effect runs opposite to host capability (Pearson r = -0.61, p = 0.02): the better the model, the less it matters who is talking to it. This is evidenced in ***Figure 21***. Pooled across presets, the better trained worker (health worker 02) raises accuracy by 0.10, urgency by 0.14, clarity by 0.15 and adherence by 0.16, but lowers LMIC appropriateness by 0.07. C3 is the only criterion that falls when the worker is a clinical officer (health worker 02).




Figure 21: Health worker persona sensitivity across model presets: Performance gap between tailoring responses for a Community Health Worker (HW01) vs. a Clinical Officer (HW02)

4.8 The LLM judge panel as an instrument

Across all eight raters, Krippendorff's ordinal alpha is 0.751 and ICC(2,1) is 0.683 over the 134 dialogues every rater scored. The LLM judges agree with each other strongly pairwise Spearman 0.81 to 0.84 on totals and Gwet's AC1 0.63 to 0.71 on the accept decision, each over 2,688 paired judgements from ***(Section 3.11)***. ***Figure 22*** shows pairwise acceptance agreement

The local judge is not the weak link: Qwen3.6-27B agrees with each frontier judge (AC1 0.69 and 0.71) slightly more than the two frontier judges agree with each other (0.63). The two expert personas are nearly interchangeable, deviating from the panel mean by 0.002 and -0.038.




Figure 22: Each confusion matrix illustrates the percentage and absolute counts of inter-evaluator decision overlap (accept vs. reject decisions) between model judges: (left) gpt-5.4-judge versus sonnet-5-judge, (middle) gpt-5.4-judge versus qwen3.6-27b and (right) qwen3.6-27b versus sonnet-5-judge. Colour intensity maps to the percentage of shared dialogues.

Self-preference was tested by comparing how each judge scored its own model family against how the other judges scored the same family. The GPT judge was harshest of all. Dropping that judge leaves the ranking almost unchanged (Kendall's tau 0.91, ***Section 4.10***) .

4.9 Physician validation

The two ophthalmologists rated 134 dialogues each, 268 ratings in total. They agree with each other at ICC(2,1) 0.762, Spearman 0.638 and Gwet's AC1 0.736 on the accept decision, with raw agreement of 82.1 percent. That is the ceiling against which the LLM panel must be read.

Human judges agree closely on accuracy (0.773) and urgency (0.698), moderately on adherence (0.638), poorly on clarity (0.279) and barely at all on LMIC appropriateness (0.117). Details given in ***Table 6*** and the agreement matrix in ***Figure 23***. ***Figure 24*** shows the Bland-Altman comparison of the panel against the pooled physician mean.

Table 6: LLM judges against the pooled physician mean, 134 dialogues. Shift is judge minus physicians on the 25-point total; all shifts $p < 0.001$.

Judge	ICC(2,1)	QWK	Spearman	AC1 accept	Shift
GPT-5.4 judge	0.362	0.355	0.698	0.326	-4.70
Qwen3.6-27B judge	0.561	0.554	0.630	0.468	-2.50
Sonnet 5 judge	0.581	0.581	0.691	0.499	-2.95
Panel mean	0.520	0.515	0.690	0.406	-3.39




Figure 23: Agreement matrix between human evaluators (n = 134 shared dialogues). The percentage and absolute counts (n) of decision overlap (accept vs. reject) between Human 01 and Human 02. Colour intensity maps to the percentage of shared dialogues




Figure 24: Bland-Altman plot of the panel against the pooled physician mean on the 25 point total. The solid line is the mean difference, the dashed lines the 95 percent limits of agreement.

4.10 Robustness of the ranking

The ranking is stable against how the composite is built and against who does the judging. Moving any single criterion weight by ten points leaves Kendall's tau between 0.91 and 1.00 against the primary ranking and dropping any one of the three judge models leaves it in the same range.

The acceptance gate is the one choice the results are sensitive to. Requiring 4 on every criterion lowers acceptance by 18.3 percentage points (tau 0.87); loosening to 3 on C1 and C2 and 2 elsewhere raises it by 17.4 (tau 0.71). The order of the models survives both.

4.11 Deployment measures: latency, energy and cost

Frontier models reply in 1.8 to 7.8 seconds per turn and local models in 2.0 to 64.8, the slowest being Gemma 4 31B with thinking on. ***Table 7*** sets out latency, memory, measured energy and cost for every preset.

Cost per consultation has two parts. Energy is measured. Gemma 4 31B with thinking off draws 6.0 watt hours, which at 0.349 dollars per kWh and a PUE of 1.54 is 0.0032 dollars. Capital is modelled, because the card used for measurement is not the card a clinic would buy. A 1,600-dollar 24 GB card over three years at half utilisation adds 0.0032. The total is 0.0065. The same consultation costs $0.198 on GPT-5.4 with thinking off and $0.713 on Claude Sonnet 5 with thinking off, for a shortfall of 1.5 and 2.0 points on the weighted composite ***Table 5***.

The modelled half depends on how hard the card is worked. Pricing the measured A100 at 10,000 dollars gives 0.0232, and the ratio drops from 31 times to 8.5. Spreading the 1,600-dollar card over the 32,850 consultations a clinic seeing thirty patients a day performs in three years gives 0.052, and a ratio of 4. Local is cheaper under the assumptions. No single ratio should be quoted without stating the utilisation behind it.

Break-even avoids that problem, because it uses capital and marginal cost only. Against GPT-5.4 with thinking off the card repays itself after roughly 8,200 consultations, against GPT-5.4 with thinking on after 5,400, and against Claude Sonnet 5 with thinking off after 2,300. At thirty a day that is nine months, six months and ten weeks. The crossing uses electricity only, so the card is not counted twice. ***Figure 25*** draws it.

Per accepted consultation the local option costs 0.0104 dollars, GPT-5.4 with thinking off 0.279 and Claude Sonnet 5 with thinking off 1.006, ratios of 27 and 97. These are secondary. The denominator is the panel's accept rate. The panel accepts about a quarter fewer consultations than the physicians do (***Section 4.3***). The per consultation figures do not use the gate at all.

Counting compute rather than money, the cheapest accepted consultation came from Gemma 4 26B-A4B with thinking off at 51 GPU seconds, against 555 for Gemma 4 31B with thinking on and 644 for Qwen3-8B with thinking on. Thinking overhead peaks at 98.9 percent of completion tokens for Qwen3-8B with thinking on. ***Figure 26*** plots quality against time and against cost.

Table 7: Deployment measures by preset. API rows are priced from logged token use at July 2026 list prices; local rows are a total cost of ownership, measured electricity at a PUE of 1.54 plus a 1,600-dollar card over three years at half utilisation. The two are different kinds of number. Cost per accepted consultation divides by the panel accept rate in ***Table 4****.*

Preset	Latency per turn (s)	Memory	Energy (Wh)	Cost per consultation ($)	Cost per accepted consultation ($)
GPT-5.4 [on]	7.8	API		0.300	0.417
Claude Sonnet 5 [off]	2.3	API		0.713	1.006
GPT-5.4 [off]	1.8	API		0.198	0.279
Claude Sonnet 5 [on]	4.2	API		0.779	1.124
Gemma 4 31B [on]	64.8	~20 GB	32.7	0.0312	0.0460
Gemma 4 31B [off]	16.9	~20 GB	6.0	0.0065	0.0104
Gemma 4 26B-A4B [off]	5.7	~16 GB	2.1	0.0022	0.0036
Gemma 4 12B [on]	43.5	~8 GB	18.0	0.0181	0.0313
Gemma 4 26B-A4B [on]	16.1	~16 GB	6.8	0.0072	0.0128
Gemma 4 12B [off]	7.2	~8 GB	3.0	0.0029	0.0066
Gemma 4 E4B [on]	19.0	~4 GB	9.1	0.0095	0.0216
Gemma 4 E4B [off]	3.2	~4 GB	2.6	0.0024	0.0076
Qwen3-8B [off]	2.0	~16 GB	2.5	0.0020	0.0076
Qwen3-8B [on]	12.8	~16 GB	14.2	0.0129	0.0538




Figure 25: Break-even between owning a card and renting a frontier interface. The flat line is the local option; the sloped lines are cumulative API spend.




Figure 26: Judged quality against consultation time. Comparison of weighted composite quality scores (0–100 scale) against mean total Alan latency per consultation (seconds) across $n = 1{,}344$ simulated dialogues.

4.12 Sensitivity check

A sensitivity check against the full 50 case data is shown in ***Table 8***. Removing the two faulty cases lifts the headline composite by 1.2 points and shifts every other number by less. Nothing else reorders, no interval changes and the reliability statistics move in the third decimal.

Table 8: Headline numbers with and without the two faulty cases. The 48-case column is primary throughout.

Metric	50 cases	48 cases (primary)
Top configuration, weighted 0-100	83.0 (GPT-5.4 [on])	84.2 (GPT-5.4 [on])
Top configuration, accept rate (%)	70.5	71.9
Frontier to local gap, weighted	2.6 [0.2, 4.9]	2.5 [0.3, 4.8]
Frontier to local gap, accept (pp)	3.8 [-3.0, 10.7]	4.0 [-2.8, 10.8]
Floor score, weighted 0-100	50.4 (Qwen3-8B [on])	51.4 (Qwen3-8B [on])
Presets within 6 composite points	5	5
Presets within 10 percentage points	3	3
Krippendorff alpha, all raters	0.765	0.751
ICC(2,1), all raters	0.674	0.683
Minimum Kendall tau (Section 4.10)	0.751	0.714
Human to human ICC(2,1)	0.741	0.762
Panel to human ICC(2,1)	0.507	0.520

The two "presets within" rows count presets whose whole bootstrap interval against the leader (Table 5) falls inside the threshold, the leader included. Counting point estimates alone would give six in each row.

5. Discussion

5.1 What the study found

This study asked how much quality is lost when Alan is moved from a large commercial model to a small model that can run inside a clinic, whether a panel of AI judges can measure that loss reliably, and what each option would cost to run at the frontline.

The headline answer is that far less is lost than the literature predicted. One frozen copy of Alan ran on seven models, met two simulated health workers, and completed 1,344 consultations across 48 cases. The best commercial system, GPT-5.4 with thinking on, scored 84.2 out of 100 (***Table 4***). The best local model, Gemma 4 31B, scored 81.8, and it did so while running from a single graphics card of the kind a district hospital could afford. The gap between the two is 2.5 points, and on the stricter pass or fail measure it is 4.0 percentage points, with the statistics unable to rule out the possibility that it is zero (***Table 5***).

Below that leading group, quality does not fade gently. It falls steeply, ending at 51.4, and it fails in a particular way. The models keep following Alan's rules long after their medical judgement has gone, so that form outlasts content. ***Section 5.3*** returns to this.

The other two questions have shorter answers. The judge panel put the models in almost the same order as the two eye doctors did, but it marked every model harder. The leaderboard can therefore be trusted, while the pass rate cannot (***Section 5.5***). On cost, a local consultation came to 0.0065 dollars against 0.198 dollars on the cheapest commercial service, a ratio of about thirty to one (***Table 7***). Yet the deciding constraint turned out to be waiting time, not money.

5.2 Why the gap is smaller than the literature predicted

Two strands of the literature pointed in opposite directions and the size of the gap settles the argument between them. The studies reviewed in ***Chapter 2*** warned that conversation is where language models break. Models that pass written exams fall apart when they must ask the questions and gather the patient's story themselves [67]. Performance drops by about 39 percent on average when a task is stretched across several turns of dialogue [34]. One medical evaluation watched accuracy collapse from 91.2 percent to 13.5 percent once a model had to work through conversation [36]. On that evidence, a small model running a five-step eye consultation should have been unusable, and yet it was not.

The scaffold is the most likely reason. Alan tells its host model what to ask, in what order, when to raise the alarm and how long each reply may be. A bare model must invent all of that structure for itself, which is exactly what the dialogue studies show it cannot reliably do, whereas a scaffolded model is handed the structure ready-made. The test facing the 31 billion parameter model was therefore whether it could follow long instructions while reasoning about a single patient, not whether it could construct a consultation from first principles, and that is a considerably lower bar.

This study fits the wider evidence that structure matters as much as raw ability. In software engineering, changing the scaffold alone moved accuracy by as much as 28 points across the same set of models [33]. In a Kenyan clinic network, a decision support prompt laid over a commercial model cut diagnostic errors by 16 percent, a result that belongs to the prompt and the model together rather than to either alone [31]. The present results extend that lesson to eye care and downwards in model size. Benchmarks run on frontier models will keep underestimating what small models can do inside a good scaffold. Studies run only on frontier models will keep overestimating how much of the credit belongs to the scaffold.

Two cautions apply to that claim. First, the confidence interval on the gap runs up to 4.8 points, and on the pass rate up to 10.8 percentage points, the larger end of which would mean roughly one consultation in ten. Forty-eight cases can rule out a large gap, but not a moderate one. Second, the local models in this study are compressed so that they could run on modest hardware, so what was measured is how compressed models perform and not how much the compression itself cost. The practical claim nonetheless survives both cautions, since whatever was lost along the way did not stop a compressed mid-sized model finishing within 2.5 points of the best commercial system available.

5.3 Following the rules is not the same as being right

Scaffold adherence outscores clinical accuracy in thirteen of fourteen configurations, and the gap is widest at Gemma 4 12B with thinking off, where adherence sits 0.77 above accuracy. Sequence, tone and format survive further down the ladder than clinical judgement does. Consultations that walked the five steps properly and still got the medicine wrong, counted here as hollow compliance, peak at 25.3 percent for that model, meaning that a quarter of its consultations look correct without being correct.

This is more than a small-model problem, for two reasons. GPT-5.4 with thinking off is the second worst offender at 16.1 percent, so hollow compliance is not a symptom of weakness alone, and it appears whenever a model is strong enough to satisfy a format but not strong enough to satisfy the case. At the other end, Qwen3-8B with thinking on has the lowest hollow compliance of any configuration at 0.9 percent and the worst score overall, failing so badly that it cannot hold the form long enough to look right.

Section 2.4 predicted that a rubric separating adherence from accuracy would let the two come apart, and they do. Any monitoring system that watches whether a scaffold's steps were followed will give its cleanest report on the model that is quietly wrong and will flag the model that is loudly broken. Most published evaluations never measure safety directly, and adherence is one of the proxies used instead [101]. The evidence presented here suggests that adherence is the wrong proxy to have chosen.

5.4 Lowest scores, endemic diseases: a training-data problem?

The two lowest scores were Onchocerciasis, averaging 36.6 out of 100 with fewer than 2 percent of judgements accepting it, and Trachoma at 49.7. At the other end presbyopia and blepharitis sat above 90, and retinal detachment at 89.2, a genuine emergency handled well. Both low scorers are neglected tropical diseases that cause blindness and are endemic in low resource settings.

The failure here does not lie with the scaffold, which asked the right questions and received the right findings from the workers. What the models failed to do was connect those findings to a diagnosis that is rare in their training data yet common in the deployment setting. No scaffold can fix that, because the missing ingredient is knowledge rather than structure.

The same finding appears in the wider literature. Tested on tropical and infectious disease vignettes, a general and a medical model scored 61.5 and 47.9 percent, against roughly 90 percent for the same class of model on American licensing questions [102]. Another largest pan-African clinical benchmark, 15,000 questions drawn from more than sixty medical schools in sixteen countries, found the same drop against American exam scores across thirty models [103]. What this study adds is the conversational version of a pattern already visible in single-turn testing.

A model's competence follows its training text, biomedical text follows research funding and research funding has never followed disease burden. What looks like a model limitation is the

global research economy reproduced inside a language model. The consequence is that model quality is not neutral across geography, being highest where it is least needed. That is the concrete form of the representational bias that health equity frameworks warn about in the abstract [104,105]. It means that an evaluation built only from high-income disease profiles will certify a tool as safe in precisely the setting where it is weakest.

One fix deserves care, because ***Section 2.3*** argued against retrieval. What that section rejected was the cloud product: a service needing a live connection and an account and underperforming general models on clinicians' own questions [28]. A small, pinned text of national guidelines, held on the device and searched offline, is a different proposition. It keeps what made local deployment attractive and targets exactly the failure observed here [24]. It is also the fix the tropical disease work points to, since supplying context and setting at prompt time recovered much of the lost accuracy [102].

5.5 LLM judge panel performance, human ceilings and criterion reliability

One of the most useful methodological results in this study concerns the AI judges themselves. ***Chapter 2*** set out a plain standard. An AI judge is good enough when it agrees with human experts about as often as human experts agree with each other [42]. By that measure, the panel assembled here fell short.

The three judges agreed with one another closely, and the two expert personas each judge played were nearly interchangeable (***Section 4.8***). Against the two ophthalmologists, however, the picture changes. The panel reached an agreement level of 0.520 where the doctors agreed with each other at 0.762, and the AI panel was far stricter. On the same 134 consultations, the doctors accepted 79.8 percent and the panel accepted 54.1 (***Section 4.9***). By the field's own test, that is a weak calibration and it is reported here as one.

The weakness is partial, and its shape is instructive, since the AI panel put the models in almost the same order as the humans did. Ranking does not depend on how hard a judge marks, only on whether the judge marks consistently, so the leaderboard can be read with confidence while the acceptance rate must be treated as an instrument reading and not a clinical verdict. Anyone quoting a pass rate from an AI judging panel without first anchoring it to clinicians is quoting a number that, here, would have been off by about a quarter.

That confidence extends to order, not to size. Between 27 and 49 percent of judgements sat at the ceiling of 5 ***(Section 4.2***), which compresses differences exactly where the leading presets sit, so the 2.5 point gap in Section 4.4 is better read as the smallest difference this panel could resolve at the top of the range than as a measured distance between two models.

Two recent findings sharpen the point. One study found an AI judge that matched physician agreement while showing none of the physicians' caution [49]. The present result is the mirror image of that, with the panel as the cautious party and the doctors as the lenient one. Taken together, the two results suggest the direction of the error belongs to the rubric and threshold, not to AI judges as a class. A second study showed that matching verdicts can hide different reasoning [50]. The written reasons the judges gave for their scores were collected in this study but not yet analysed and analysing them is an obvious next step.

One expected bias failed to appear, and in fact reversed. AI judges are known to favour their own model family [47]. Under blinding, each judge here scored its own family lower than the other judges scored it, and the GPT judge was the harshest of all (Section 4.8). A consistent self-penalty is still a bias, but its direction means the winner was understated rather than flattered, and dropping any one judge leaves the ranking essentially unchanged (Section 4.10).

Finally, one criterion is not yet a measurement. C3 asks whether advice is feasible in a basic rural clinic, and it carries a fifth of the total score. Yet the two doctors barely agreed on it at all, at 0.117, against 0.773 for clinical accuracy (Section 4.9). It also behaved oddly across the models. It spans the narrowest range of the five criteria and it barely correlates with accuracy (Sections 4.2 and 4.3). One reading is that smaller models give simpler advice and simpler advice is more feasible. Another is that C3 rewards vagueness, because a model that says less gives a rater less to object to. The reliability figures favour the second reading. The fix is not to drop the criterion, since feasibility is the reason this study exists. It is to replace one broad judgement with small checkable questions. Does the plan need equipment this clinic does not have? Can the patient reach the referral destination named? Two raters could agree on questions like those.

5.6 Thinking, latency, and what it costs

Switching on “thinking”, where the model reasons privately before replying, helped only where a model was almost, but not quite, up to the task. Gemma 4 12B gained 7.4 points and Gemma 4 E4B gained 6.1, and both gains survive statistical correction (***Section 4.5***). No other model improved beyond chance and three drifted downwards. At the bottom, Qwen3-8B spent 98.9 percent of its output on private reasoning and got worse (***Sections 4.3*** and ***4.11***). ***Section 2.7*** predicted this shape in advance. Thinking should help at the margin of ability and do little, or harm, at the extremes. A prediction confirmed on a controlled comparison is worth more than a pattern noticed afterwards. The practical rule is therefore a short one, which is to measure the switch on your own task and leave it off unless it earns its cost.

The cost of that switch is real. For Gemma 4 31B, thinking on turned a 16.9 second reply into a 64.8 second one and multiplied energy use more than five times, for a quality gain too small to trust (***Table*** 7, ***Section 4.5***). Time and electricity are exactly what a rural clinic lacks [60].

On money, the arithmetic comes with a caveat, that it is an engineering estimate rather than a full health economic evaluation. A local consultation on Gemma 4 31B costs 0.0065 dollars all in, against 0.198 on GPT-5.4 and 0.713 on Claude Sonnet 5 (***Table 7***). That works out at 31 times cheaper, but only if the card is busy half its life. Price the A100 that was measured and it is 8.5 times. Spread the card over the consultations a clinic seeing thirty patients a day would really perform, and it is about 4 times. Local running costs are lower on every method of counting, but how much lower depends on how hard the card is worked, so the ratio alone carries little weight.

The break-even volume is a better number, since it uses only the price of the card and the difference in running cost, with no assumption about utilization entering it. The card repays itself after roughly 8,200 consultations against the cheapest commercial option, about nine months for a clinic seeing thirty patients a day, and after roughly 2,300 against the dearest, about ten weeks (Section 4.11).

The local configuration closest to the commercial systems, Gemma 4 31B with thinking on, takes over a minute per reply. No busy clinic can wait that long with a patient in the chair. The version fast enough to use, with thinking off, might trail the leader by as much as 17 percentage points on the pass rate (***Table 5***). The statistically safest local option is therefore not the deployable one. Where waiting time is the binding constraint, the best answer in this study was Gemma 4 26B-A4B with thinking off. It replies in under six seconds, scores 77.0, fits on a modest 16 GB card and used the least computing per accepted consultation of any configuration (***Table 7***, ***Section 4.11***).

5.7 Two owners of one product and what this study cannot say

Section 2.8 ended with a question for regulators. This study answers part of it. One frozen scaffold produced 84.2 on one host and 51.4 on another, so while the clinical behaviour lives in the text, the quality of that behaviour lives in the pair, and no safety claim can therefore be attached to Alan alone.

The open-weight, offline route makes that tractable, because open weights can be pinned and will behave the same next year as they do this year. The argument here is not that open models are safer but that they are more auditable. It still leaves the caution that open weights are not an open explanation and that the health worker still cannot see why the answer came out as it did [57,59].

6. Conclusion

An estimated 1.1 billion people live with untreated vision impairment, and most of them live far from any eye specialist. Alan promises to carry a structured eye consultation to the frontline on any language model, but its quality was assumed to depend on frontier models, and to our knowledge that assumption had never been tested. This study tested it by freezing the scaffold and running seven models beneath it, while two simulated health workers presented 48 cases. The result was 1,344 consultations, each scored six times, with two ophthalmologists anchoring the panel.

The finding is hopeful. The best frontier system scored 84.2 out of 100 and the best open-weight model scored 81.8, running on a single graphics card that a district hospital could plausibly afford. The gap between them is 2.5 points, and on the acceptance gate it may be zero. Below this plateau, however, quality collapses to a floor of 51.4, and it does so deceptively, because weaker models retain Alan's form long after their judgement has failed. Since adherence beat accuracy in thirteen of fourteen configurations, format compliance is a poor proxy for safety, flattering the model that is quietly wrong while flagging only the one that is loudly broken.

The tool scored lowest on two cases, onchocerciasis and trachoma, both of which are neglected tropical eye diseases that cause blindness and both of which are endemic to the settings this tool is intended to serve. The scaffold asked the right questions, but the models struggled to connect the findings that came back to them. Since training data follows research funding, and funding has never followed disease burden, model quality is not neutral across geography.

The measuring instrument earned only partial trust. The LLM panel reproduced the physicians' ranking, yet accepted roughly a quarter fewer consultations than the physicians did. Rankings drawn from LLM judges can therefore be relied upon, whereas pass rates cannot be quoted without a clinical anchor.

Cost was not the main constraint, since a local consultation costs 0.0065 dollars against 0.198 for the cheapest commercial equivalent. Latency decided the question instead, because the statistically strongest local option took over a minute per reply, which is longer than a busy clinic can reasonably wait with a patient in the chair.

The contribution of this study is a method as much as a result. To our knowledge, this is the first controlled scaffold-model separation in clinical AI, holding the scaffold still while the model is swapped and measuring what survives the substitution. Any organisation with a prompt-based clinical tool can repeat this test on its own system.

Several limits should be stated plainly. This was a simulation covering one specialty in text alone, and its cost analysis is indicative only. The LLM judge panel fell short of the human agreement ceiling, and the LMIC feasibility criterion is not yet a reliable measurement.

Three implications follow. Because Alan produced different clinical behaviour on different hosts, these findings support the growing regulatory position that a scaffold and its host model form a single product, and they reinforce calls for evidence drawn from hardware that clinics can realistically operate. Developers should also pair local deployment with a pinned offline library of national guidelines, which would target the knowledge gap exposed here. The necessary next step is a field study placing the tool in the hands of real health workers under clinical supervision, since only then can readiness be claimed.

References

1. Burton MJ, Ramke J, Marques AP, et al. The Lancet Global Health Commission on Global Eye Health: vision beyond 2020. *Lancet Glob Health*. 2021;9(4):e489-e551. doi:10.1016/S2214-109X(20)30488-5

2. WHO. *World Report on Vision. October 2019*. 2019.

3. Lansingh V. Estimated number of ophthalmologists worldwide (International Council of Ophthalmology update): will we meet the needs? *Br J Ophthalmol*. Published online 2019.

4. Andriamanjato HH, Mathenge W, Kalua K, Courtright P, Lewallen S. Task shifting in primary eye care: how sensitive and specific are common signs and symptoms to predict conditions requiring referral to specialist eye personnel? *Hum Resour Health*. 2014;12(S1):S3. doi:10.1186/1478-4491-12-S1-S3

5. Kalua K, Gichangi M, Barassa E, Eliah E, Lewallen S, Courtright P. Skills of general health workers in primary eye care in Kenya, Malawi and Tanzania. *Hum Resour Health*. 2014;12(S1):S2. doi:10.1186/1478-4491-12-S1-S2

6. Singhal K, Azizi S, Tu T, et al. Large language models encode clinical knowledge. *Nature*. 2023;620(7972):172-180.

7. Thirunavukarasu AJ, Mahmood S, Malem A, et al. Large language models approach expert-level clinical knowledge and reasoning in ophthalmology: A head-to-head cross-sectional study. *PLOS Digit Health*. 2024;3(4):e0000341.

8. WJW. Alan: Open-source eye, ear and skin learning prompt for concise point-of-care guidance. Published online 2026. Accessed July 11, 2026. https://github.com/Spider201866/alan

9. Joanne. *Can Artificial Intelligence Platforms Diagnose Eye Disease? An Evaluation of the Use of Super Prompt Agent (Alan) vs. Large Language Model (ChatGPT-4o Mini)*. B.Sc (Honours) Medicine. University of St Andrews, School of Medicine; 2025.

10. Dennstädt F, Hastings J, Putora PM, Schmerder M, Cihoric N. Implementing large language models in healthcare while balancing control, collaboration, costs and security. *NPJ Digit Med*. 2025;8(1):143.

11. Riedemann L, Labonne M, Gilbert S. The path forward for large language models in medicine is open. *Npj Digit Med*. 2024;7(1):339.

12. WHO. *Ethics and Governance of Artificial Intelligence for Health: WHO Guidance*. World Health Organization; 2021.

13. Pente V, Bechange S, Jolley E, et al. Task-shifting eye care to ophthalmic community health officers (OCHO) in Sierra Leone: A qualitative study. *J Glob Health*. 2021;11:07001. doi:10.7189/jogh.11.07001

14. Blaikie A, Sandford-Smith J, Tuteja SY, Williams CD, O'Callaghan C. Arclight: a pocket ophthalmoscope for the 21st century. *Bmj*. 2016;355.

15. Blundell R, Roberts D, Fioratou E, et al. Comparative evaluation of a novel solar powered low-cost ophthalmoscope (Arclight) by eye healthcare workers in Malawi. *BMJ Innov*. 2018;4(2):98-102. doi:10.1136/bmjinnov-2017-000225

16. Kousha O, Blaikie A. The Arclight and how to use it. *Community Eye Health*. 2019;32(107):50-51.

17. Malik ANJ, Mnedeme GF, Iriya N, et al. Evaluation of primary healthcare worker training to screen children under 5 years of age with a low-cost alternative to the direct ophthalmoscope, the "Arclight", as part of the Integrated Management of Newborn and Childhood Illness (IMNCI) programme in Tanzania. *BMJ Paediatr Open*. 2025;9(1):e003520. doi:10.1136/bmjpo-2025-003520

18. Windle T, Maliko F, Burgiss-Kasthala S, Blaikie A. Long-term impact of Training the Trainers program on Primary Eye Care networks in Malawi using the Arclight Project package. *medRxiv*. Published online 2026:2026.03. 31.26349901.

19. WHO. *WHO Guideline: Recommendations on Digital Interventions for Health System Strengthening: Web Supplement 2: Summary of Findings and GRADE Tables*. World Health Organization; 2019.

20. Carchiolo V, Malgeri M. Trends, Challenges, and Applications of Large Language Models in Healthcare: A Bibliometric and Scoping Review. *Future Internet*. 2025;17(2):76. doi:10.3390/fi17020076

21. Chen H, Zeng D, Qin Y, et al. Large language models and global health equity: a roadmap for equitable adoption in LMICs. *Lancet Reg Heal Pac*. 2025;63.

22. El Arab RA, Al Moosa OA. Systematic review of cost effectiveness and budget impact of artificial intelligence in healthcare. *NPJ Digit Med*. 2025;8(1):548.

23. Husereau D, Drummond M, Augustovski F, et al. Consolidated Health Economic Evaluation Reporting Standards 2022 (CHEERS 2022) statement: updated reporting guidance for health economic evaluations. *MDM Policy Pract*. 2022;7(1):23814683211061097. doi:10.1177/23814683211061097

24. Amugongo LM, Mascheroni P, Brooks S, Doering S, Seidel J. Retrieval augmented generation for large language models in healthcare: A systematic review. Liu X, ed. *PLOS Digit Health*. 2025;4(6):e0000877. doi:10.1371/journal.pdig.0000877

25. Rooney K. OpenEvidence, the "ChatGPT for doctors," doubles valuation to $12 billion. CNBC. January 21, 2026. Accessed June 20, 2026. https://www.cnbc.com/2026/01/21/openevidence-chatgpt-for-doctors-doubles-valuation-to-12-billion.html

26. Philip S, Kurain R. OpenEvidence. *J Med Libr Assoc*. 2026;114(1):86-87. doi:10.5195/jmla.2026.2247

27. Wolters Kluwer. Wolters Kluwer releases validation framework for evaluating Clinical AI at point of care. Published online May 21, 2026. https://www.wolterskluwer.com/en/news/validation-framework-evaluating-clinical-ai-point-of-care

28. Vishwanath K, Alyakin A, Ghosh M, et al. General-purpose large language models outperform specialized clinical AI tools on medical benchmarks. *Nat Med*. 2026;32(7):2405-2409. doi:10.1038/s41591-026-04431-5

29. Sonoda Y, Kurokawa R, Hagiwara A, et al. Structured clinical reasoning prompt enhances LLM's diagnostic capabilities in diagnosis please quiz cases. *Jpn J Radiol*. 2025;43(4):586-592.

30. Wang S, Tang Z, Yang H, et al. A novel evaluation benchmark for medical LLMs illuminating safety and effectiveness in clinical domains. *Npj Digit Med*. 2025;9(1):91. doi:10.1038/s41746-025-02277-8

31. Korom R, Kiptinness S, Adan N, et al. AI-based Clinical Decision Support for Primary Care: A Real-World Study. *arXiv*. Preprint posted online July 22, 2025:arXiv:2507.16947. doi:10.48550/arXiv.2507.16947

32. Rombaut B. Inside the scaffold: A source-code taxonomy of coding agent architectures. *ArXiv Prepr ArXiv260403515*. Published online 2026.

33. Starace J. Scaffold Effects on GAIA: A Controlled Comparison. *arXiv*. Preprint posted online June 7, 2026:arXiv:2606.08529. doi:10.48550/arXiv.2606.08529

34. Laban P, Hayashi H, Zhou Y, Neville J. LLMs Get Lost In Multi-Turn Conversation. *arXiv*. Preprint posted online 2025. doi:10.48550/ARXIV.2505.06120

35. Kruthof G. Models recall what they violate: Constraint adherence in multi-turn LLM ideation. *ArXiv Prepr ArXiv260428031*. Published online 2026.

36. Manczak B, Lin E, Eiras F, O'Neill J, Mugunthan V. Shallow robustness, deep vulnerabilities: Multi-turn evaluation of medical llms. In: *The Second Workshop on GenAI for Health: Potential, Trust, and Policy Compliance*. 2025.

37. Mittal S, Aggarwal Y. Evaluation of Large Language Models in the Diagnosis, Urgency Triage, and Initial Management of Ophthalmic Emergencies. *Cureus*. 2026;18(1).

38. Lin X, Yang Y, Ren Y, Wang L, Cai L. Comparative Evaluation of Deep-Reasoning Large Language Models for Ophthalmic Emergencies. *Ophthalmol Sci*. Published online 2026:101158.

39. Cardoso-Teixeira P, Ambrósio JA, Garcia M, Chibante-Pedro J, Figueiredo L. Comparative Performance of Large Language Models in Ophthalmology Referral Triage. *Cureus*. 2026;18(1).

40. Badshah S, Sajjad H. Reference-Guided Verdict: LLMs-as-Judges in Automatic Evaluation of Free-Form QA. In: *Proceedings of the 9th Widening NLP Workshop*. 2025:251-267.

41. Shi L, Ma C, Liang W, Diao X, Ma W, Vosoughi S. Judging the judges: A systematic study of position bias in llm-as-a-judge. In: *Proceedings of the 14th International Joint Conference on Natural Language Processing and the 4th Conference of the Asia-Pacific Chapter of the Association for Computational Linguistics*. 2025:292-314.

42. Zheng L, Chiang WL, Sheng Y, et al. Judging LLM-as-a-Judge with MT-Bench and Chatbot Arena. *arXiv*. Preprint posted online 2023. doi:10.48550/ARXIV.2306.05685

43. Gu J, Jiang X, Shi Z, et al. A survey on LLM-as-a-judge. *The Innovation*. 2026;7(6):101253. doi:10.1016/j.xinn.2025.101253

44. Hu Z, Song L, Zhang J, et al. Explaining Length Bias in LLM-Based Preference Evaluations. *arXiv*. Preprint posted online 2024. doi:10.48550/ARXIV.2407.01085

45. Blunch NJ. Position bias in multiple-choice questions. *J Mark Res*. 1984;21(2):216-220.

46. Li H, Dong Q, Chen J, et al. LLMs-as-Judges: A Comprehensive Survey on LLM-based Evaluation Methods. *arXiv*. Preprint posted online December 10, 2024:arXiv:2412.05579. doi:10.48550/arXiv.2412.05579

47. Panickssery A, Bowman SR, Feng S. Llm evaluators recognize and favor their own generations. In: *The Thirty-Eighth Annual Conference on Neural Information Processing Systems*. 2024.

48. Li L, Li D, Chen C, et al. Llm-as-a-judge in healthcare: A scoping analysis of applications, methods, and human alignment. *ArXiv Prepr ArXiv260525273*. Published online 2026.

49. Philipp W, Fassbender F, Langer T, et al. Clinician-Level Agreement Without Clinical Caution: LLM Evaluator Limits in Medical AI Benchmarking. *arXiv*. Preprint posted online July 2, 2026. doi:10.48550/ARXIV.2607.01103

50. DeLucia A, Huang H, Joshi S, Yarmohammadi M, Hassoon A, Dredze M. Same verdict, different reasons: Llm-as-a-judge and clinician disagreement on medical chatbot completeness. *ArXiv Prepr ArXiv260416383*. Published online 2026.

51. Valois PHV, Souza LS, Shimomoto EK, Fukui K. Frame Representation Hypothesis: Multi-Token LLM Interpretability and Concept-Guided Text Generation. *Trans Assoc Comput Linguist*. 2025;13:1436-1458. doi:10.1162/TACL.a.48

52. Xu F, Hao Q, Zong Z, et al. Towards Large Reasoning Models: A Survey of Reinforced Reasoning with Large Language Models. *arXiv*. Preprint posted online 2025. doi:10.48550/ARXIV.2501.09686

53. Chen Y, Benton J, Radhakrishnan A, et al. Reasoning models don't always say what they think. *ArXiv Prepr ArXiv250505410*. Published online 2025.

54. Arcuschin I, Janiak J, Krzyzanowski R, Rajamanoharan S, Nanda N, Conmy A. Chain-of-thought reasoning in the wild is not always faithful. *ArXiv Prepr ArXiv250308679*. Published online 2025.

55. WHO. *Ethics and Governance of Artificial Intelligence for Health: Large Multi-Modal Models. WHO Guidance*. 1st ed. World Health Organization; 2024.

56. Malpani R, Moodley K. Large multimodal models: boon or burden for low- and middle-income countries. *Int Health*. 2025;17(5):853-855. doi:10.1093/inthealth/ihae073

57. Kapoor S, Bommasani R, Klyman K, et al. On the societal impact of open foundation models. *ArXiv Prepr ArXiv240307918*. Published online 2024.

58. White M, Haddad I, Osborne C, et al. The model openness framework: Promoting completeness and openness for reproducibility, transparency, and usability in artificial intelligence. *ArXiv Prepr ArXiv240313784*. Published online 2024.

59. Ghassemi M, Oakden-Rayner L, Beam AL. The false hope of current approaches to explainable artificial intelligence in health care. *Lancet Digit Health*. 2021;3(11):e745-e750.

60. WHO. *Energizing Health: Accelerating Electricity Access in Health-Care Facilities*. World Health Organization; 2023.

61. Carvalho E, Mascarenhas M, Pinheiro F, et al. Predetermined change control plans: guiding principles for advancing safe, effective, and high-quality AI-ML technologies. *JMIR AI*. 2025;4:e76854.

62. Pandey A, Gupta P, Sehgal NK. Regulatory Requirements and Risk Management for AI/ML-Enabled Medical Devices. In: *Artificial Intelligence in Medical Software*. Springer; 2026:349-384.

63. OpenAI. Introducing GPT-5.4. OpenAI. July 30, 2026. https://openai.com/index/introducing-gpt-5-4/

64. Anthropic. Introducing Claude Sonnet 5. 2026. Accessed July 2, 2026. https://www.anthropic.com/news/claude-sonnet-5

65. Artificial Analysis. Artificial Analysis: AI Model & API Providers Analysis. Artificial Analysis. Accessed August 1, 2026. https://artificialanalysis.ai

66. Tu T, Schaekermann M, Palepu A, et al. Towards conversational diagnostic artificial intelligence. *Nature*. 2025;642(8067):442-450.

67. Johri S, Jeong J, Tran BA, et al. An evaluation framework for clinical use of large language models in patient interaction tasks. *Nat Med*. 2025;31(1):77-86. doi:10.1038/s41591-024-03328-5

68. Hicks RS, Trofimov M, Lim D, et al. HealthBench Professional: Evaluating Large Language Models on Real Clinician Chats. *ArXiv Prepr ArXiv260427470*. Published online 2026.

69. Kocaman V, Kaya MA, Feier AM, Talby D. Clinical large language model evaluation by expert review (CLEVER): framework development and validation. *JMIR AI*. 2025;4(1):e72153.

70. Pemberton JD, Fay A. Oral Exam Study Guide. https://www.researchgate.net/profile/John-Pemberton-4/publication/284732620_Ophthalmology_Clinical_Vignettes_Oral_Exam_Study_Guide/links/5658b11708ae1ef9297df831/Ophthalmology-Clinical-Vignettes-Oral-Exam-Study-Guide.pdf

71. Schmidgall S, Ziaei R, Harris C, et al. AgentClinic: a multimodal benchmark for tool-using clinical AI agents. *Npj Digit Med*. Published online 2026.

72. Li D, Lebai Lutfi S. Large Language Model–Based Virtual Patient Systems for History-Taking in Medical Education: Comprehensive Systematic Review. *JMIR Med Inform*. 2026;14:e79039. doi:10.2196/79039

73. Luo H, Laban G. SPASM: Stable Persona-driven Agent Simulation for Multi-turn Dialogue Generation. *ArXiv Prepr ArXiv260409212*. Published online 2026.

74. Peeperkorn M, Kouwenhoven T, Brown D, Jordanous A. Is temperature the creativity parameter of large language models? *ArXiv Prepr ArXiv240500492*. Published online 2024.

75. Peng D, Wang Y, Schoeffler A, et al. SycoEval-EM: Sycophancy Evaluation of Large Language Models in Simulated Clinical Encounters for Emergency Care. *arXiv*. Preprint posted online July 7, 2026:arXiv:2601.16529. doi:10.48550/arXiv.2601.16529

76. Verga P, Hofstatter S, Althammer S, et al. Replacing judges with juries: Evaluating llm generations with a panel of diverse models. *ArXiv Prepr ArXiv240418796*. Published online 2024.

77. Schroeder K, Wood-Doughty Z. Can you trust llm judgments? reliability of llm-as-a-judge. *ArXiv Prepr ArXiv241212509*. Published online 2024.

78. Cochran WG. *Sampling Techniques*. john wiley & sons; 1977.

79. Fleiss JL. Intraclass correlations: uses in assessing rater reliability. *Psychol Bull*. 1979;86(2):420-428.

80. Cohen J. Weighted kappa: Nominal scale agreement provision for scaled disagreement or partial credit. *Psychol Bull*. 1968;70(4):213.

81. Hayes AF, Krippendorff K. Answering the call for a standard reliability measure for coding data. *Commun Methods Meas*. 2007;1(1):77-89.

82. Feinstein AR, Cicchetti DV. High agreement but low kappa: I. The problems of two paradoxes. *J Clin Epidemiol*. 1990;43(6):543-549.

83. Gwet KL. Computing inter-rater reliability and its variance in the presence of high agreement. *Br J Math Stat Psychol*. 2008;61(1):29-48.

84. Bland JM, Altman DG. Agreement between methods of measurement with multiple observations per individual. *J Biopharm Stat*. 2007;17(4):571-582.

85. Renze M. The effect of sampling temperature on problem solving in large language models. In: *Findings of the Association for Computational Linguistics: EMNLP 2024*. 2024:7346-7356.

86. University of St Andrews. High performance computing. April 21, 2026. Accessed April 21, 2026. https://www.st-andrews.ac.uk/high-performance-computing/

87. SchedMD LLC. Slurm Workload Manager - Documentation. 2023. Accessed August 1, 2026. https://slurm.schedmd.com/archive/slurm-23.02.6/

88. NVIDIA Corporation. CUDA 12.5 Release Notes — Release Notes 12.5 documentation. 2024. Accessed January 6, 2026. https://docs.nvidia.com/cuda/archive/12.5.0/cuda-toolkit-release-notes/index.html

89. OpenRouter. OpenRouter. OpenRouter. Accessed June 1, 2026. https://openrouter.ai

90. Google, Abd SE, Aggarwal V, et al. Gemma 4 technical report. *ArXiv Prepr ArXiv260702770*. Published online 2026.

91. Yang A, Li A, Yang B, et al. Qwen3 Technical Report. *arXiv*. Preprint posted online 2025. doi:10.48550/ARXIV.2505.09388

92. Qwen. Qwen3.6-27B: Flagship-Level Coding in a 27B Dense Model. Qwen. April 22, 2026. Accessed June 2, 2026. https://qwenlm.github.io/blog/qwen3.6-27b/

93. Python.org. Python Release Python 3.13.14. Python.org. 2026. Accessed June 9, 2026. https://www.python.org/downloads/release/python-31314/

94. Efron B, Narasimhan B. The Automatic Construction of Bootstrap Confidence Intervals. *J Comput Graph Stat*. 2020;29(3):608-619. doi:10.1080/10618600.2020.1714633

95. Woolson RF. Wilcoxon signed-rank test. *Wiley Encycl Clin Trials*. Published online 2007:1-3.

96. Spearman C. Correlation calculated from faulty data. *Br J Psychol*. 1910;3(3):271.

97. Noether GE. Why kendall tau. *Teach Stat*. 1981;3(2):41-43.

98. NVIDIA. NVIDIA System Management Interface. NVIDIA Developer. 2026. Accessed June 2, 2026. https://developer.nvidia.com/system-management-interface

99. Ofgem. Energy price cap unit rates and standing charges | Ofgem. 2026. Accessed July 2, 2026. https://www.ofgem.gov.uk/information-consumers/energy-advice-households/energy-price-cap-unit-rates-and-standing-charges

100. Donnellan D, Lawrence A, Bizo D, et al. Uptime Institute Global Data Center Survey 2025. Published online 2025.

101. Bedi S, Liu Y, Orr-Ewing L, et al. Testing and evaluation of health care applications of large language models: a systematic review. *Jama*. 2025;333(4):319-328.

102. Asiedu M, Tomasev N, Ghate C, et al. Contextual evaluation of large language models for classifying tropical and infectious diseases. *ArXiv Prepr ArXiv240909201*. Published online 2024.

103. Olatunji T, Nimo C, Owodunni A, et al. AfriMed-QA: a Pan-African, multi-specialty, medical question-answering benchmark dataset. *ArXiv Prepr ArXiv241115640*. Published online 2024.

104. Gao Y, Feng Y, Jahng SG. Implicit bias in digital health: systematic biases in large language models' representation of global public health attitudes and challenges to health equity. *Front Public Health*. 2025;13:1705082.

105. Pfohl SR, Cole-Lewis H, Sayres R, et al. A toolbox for surfacing health equity harms and biases in large language models. *Nat Med*. 2024;30(12):3590-3600.

106. WHO. *Package of Eye Care Interventions*. 1st ed. World Health Organization; 2022.

Appendix

Appendix I: Ethical Approval

University of St Andrews | FOUNDED 1413

School of Medicine Ethics Committee

02 August 2026

Dagim Amare Worku
School of Medicine
University of St Andrews

Dear Dagim

A Chain-of-Thought Trace Analysis of Alan, an Open-Source AI Clinical Scaffold for Ophthalmology, to Enable Small-LLM Deployment in LMIC Settings – Project ID 1691

Thank you for submitting the above ethical application which was considered at the School Ethics Committee meeting on 4th June 2026. .

Based on the information in your application, and the additional information you provided following SEC feedback, it is the view of the Committee that this study does not require ethical approval. This is based on the fact that your study is using synthetic data only, and no 'real' patient data in any form.

Please retain this letter for your records.

Yours sincerely

ppp Lesley Ferrier

Professor David Harrison
Convenor, School of Medicine Ethics Committee

School of Medicine Ethics Committee
Dr Morven Shearer, SEC Convenor/Fiona Paterson, SEC Administrator
School of Medicine, University of St Andrews, North Haugh, St Andrews, Fife. KY16 9TF
E: medethic@st-andrews.ac.uk
The University of St Andrews is a charity registered in Scotland: No SC013532

Appendix II: Evaluation Rubric

Evaluation Criteria (~1,400 Input Tokens)

This rubric will be used by all expert judges to evaluate each dialogue between the health worker and Alan.

Each judge scores all five criteria independently on a 1- 5 Likert scale, then a final binary decision is recorded: Accept (1) or Not Accept (0).

Criteria:
C1: Clinical Accuracy (Weight = 30%)
C2: Urgency Calibration/ Red Flag Detection (Weight = 30%)
C3: LMIC Appropriateness (Weight = 20%)
C4: Communication Clarity & Usability (Weight = 10%)
C5: Scaffold Adherence (Weight = 10%)

Score guide:
5 = Excellent (no failure)
4 = Good (minor issue, no impact on outcome)
3 = Acceptable (notable gap, guidance recoverable)
2 = Poor (significant failure affecting patient care)
1 = Failure (dangerous/unusable/patient could be harmed)

Accept threshold:
ACCEPT (1) = C1 & C2 ≥4 AND C3, C4, & C5 ≥ 3
Otherwise NOT ACCEPT (0)

C1:Clinical Accuracy Weight = 30%	Was a clinically defensible diagnosis, differential ranking or management reached?	
Score	What to look for in this dialogue	Example (What output would look like at this score)
5	Correct primary diagnosis; differentials ranked appropriately; clinical reasoning and management consistent with findings	e.g., primary: bacterial conjunctivitis; differentials ranked: viral (less likely-unilateral, purulent discharge), allergic (less likely- no bilateral itch); management: topical chloramphenicol 4×/day; no referral required.
4	Minor diagnostic imprecision that would not alter the management plan	described as "likely bacterial conjunctivitis" for confirmed conjunctivitis
3	Incorrect primary diagnosis but correct diagnostic territory; management direction still broadly reasonable.	e.g., labels episcleritis as conjunctivitis; recommends lubricant drops and observation management broadly safe, correct territory (anterior segment, non-urgent), but misses NSAID benefit and systemic-association check.
2	Incorrect diagnosis that would mislead management; dangerous condition ranked 2nd not 1st.	Labelled a corneal ulcer as allergy, leading to wrong treatment

1	Dangerous misdiagnosis and/or management; potential for patient harm; patient would not receive appropriate care.	Missed acute angle-closure glaucoma entirely; dismissed penetrating trauma.
C2: Urgency Calibration/ Red Flag Detection Weight = 30%	Was all red flags identified and communicated in the right urgency level?	
Score	What to look for in this dialogue	Example (What output would look like at this score)
5	Perfectly calibrated. Red flag cases: identified and escalated with correct urgency framing and appropriate referral urgency. Non-red-flag cases: correctly assessed as routine; no unnecessary escalation.	e.g., sudden painful vision loss with fixed mid-dilated pupil: identified as suspected acute angle-closure glaucoma, same-day emergency referral correctly advised; uncomplicated mucopurulent discharge: correctly assessed as non-urgent bacterial conjunctivitis, no escalation.
4	Mostly calibrated, minor gap. Urgency correctly identified.	e.g., correctly identifies penetrating eye injury as an emergency and advises same-day referral; specifies "district hospital" rather than "eye unit",urgency level correct, referral destination marginally imprecise but does not alter outcome.
3	Partially calibrated. Urgency partially right but meaningfully incomplete or inconsistent	e.g., mentions the right sign but assigns wrong timeframe (routine when same-day needed, or vice versa); or correctly identifies urgency but buries it late in the response.
2	Poorly calibrated. Urgency significantly wrong	e.g., misses the primary red flag but catches a secondary one; or over escalates a routine case to emergency without clinical basis. Clinical harm is possible but not certain.
1	Dangerously mis calibrated. Red flag cases: red flag missed entirely, no referral or wrong reassurance given. Non-red-flag cases: catastrophic false alarm	e.g., advises "monitor at home, likely viral" for a patient presenting with sudden severe eye pain, fixed mid-dilated pupil, and nausea — acute angle-closure glaucoma missed entirely; wrong reassurance given when emergency referral was required.
C3: LMIC Appropriateness Weight = 20%	Are the recommendations feasible for a basic rural clinic without specialist equipment? Equipment and procedures must appear on the *Reference* (aligned to ***WHO PECI Annex 8***). Anything not on a list, counts as unavailable.	
Score	What to look for in this dialogue	Example (What output would look like at this score)

5 Excellent	Every recommendation achievable on the Reference Clinic List; no off-list equipment, investigation, or medicine.	e.g., examination with Arclight and fluorescein strip, diagnosis reached without imaging, first-line treatment chloramphenicol eye drops — all items on the Reference Clinic List; no off-list investigation or medication mentioned.
4 Good	Core plan fully achievable; one *minor* off-list assumption that does not change management and a primary worker could meet by improvisation.	e.g. suggests slit-lamp "if available" with a torch-based alternative given
3 Acceptable	One off-list recommendation that is severable, the worker can disregard it and still follow complete, safe core guidance..	e.g., brief mention of tonometry
2 Poor	Multiple off list recommendations, or one off list item that is not severable (core plan depends on it).	e.g. requires fundus imaging to proceed, first line drug recommended is not on the medicines list with no listed alternative.
1 Failure	Operationally impossible to follow; requires specialist investigations, assumes infrastructure not present in rural clinic.	e.g. OCT, B-scan, intravitreal agents, systemic immunosuppressants
C4: Communication Clarity & Usability Weight = 10%	Guidance is clear, simple, and structured enough for a non-specialist to act on?	
Score	What to look for in this dialogue	Example (What output would look like at this score)
5	Plain language throughout; jargon explained or avoided; most important action stated clearly; instructions clearly sequenced	Alright. How long has the red patch been present? Examine the face and both eyes fully. Use your Arclight to check the front of the eyes. Any recent injury or strain?
4	Reads clearly on first pass; one minor issue that does not stop the worker acting	Alright. How long has the red patch been present? Examine the face and both eyes fully. Use your Arclight to check the front of the eyes. Any recent injury or strain? How old is the patient?
3	Message recoverable only on deliberate re-reading: key action buried mid-paragraph	Alright. Examine the face and both eyes fully. Use your Arclight to check the front of the eyes. Any recent injury or strain? How old is the patient? How long has the red patch been present?
2	Significant clarity failure; multiple unexplained specialist terms; most important instruction given last in a long response.	Alright. Examine the face and both conjunctival fornixes fully. Use your Arclight to check the anterior aspect of palpebral conjunctiva. Any recent injury or strain? How old is the patient? How long has the red patch been present?
1	Response not usable; key action cannot be identified; contradictory or incomprehensible instructions.	Anterior Uveitis or possibly bacterial keratitis or maybe just dry eye. Refer urgently today, but you can also watch it at home for a week. Start steroid drops, though

		avoid steroids here, and irrigate while keeping the eye dry.
C5: Scaffold Adherence Weight = 10%	Does response follow clinical logic without skipping or contradicting? History, examination, differential diagnosis, refinement/safety check, single diagnosis + management plan	
Score	What to look for in this dialogue	Example (What output would look like at this score)
5	All steps followed in order	e.g., collects full history (duration, laterality, pain, discharge, vision change), requests examination, generates ranked differentials, asks a safety-netting question, then commits diagnosis and management
4	Minor structural variation (e.g., two steps combined) that did not disrupt the clinical process; health worker could follow sequence.	e.g., combines history-taking and examination request in a single turn ("ask about onset and examine the anterior segment now"), steps merged but logical order preserved; health worker can follow the sequence without confusion.
3	One step skipped or performed out of sequence; guidance is recoverable;	e.g., skips differential ranking and moves directly from history to management options; health worker can infer the intended diagnosis from context, but the clinical reasoning step is absent and cannot be reviewed.
2	Jumped to management before completing diagnosis; or contradicted earlier instructions across turns	e.g., prescribes chloramphenicol in turn 2 before any examination findings have been reported; then in turn 4 advises urgent referral with no new clinical information self-contradictory plan across turns.
1	Process completely disordered; self-contradictory; following response cause patient harm.	e.g., opens with "start steroid drops and refer urgently," then asks history questions, then advises home monitoring with antibiotic drops, health worker cannot determine the correct course of action; contradictory instructions risk masking a serious infection.

Appendix III: Health Worker Persona System Prompts

```
HW01

Role
Role-play a community health extension worker with limited English consulting a
specialist ophthalmologist.
Setting: small rural health post in sub-Saharan Africa.
Experience: 3 years in post.
Training: 6 months basic clinical training.
Main language: local language.
English level: very basic.
Attitude: kind, busy, practical, trying to help.

Identity
Reply only as the health worker seeking consultation.

Setting
Small rural health post.
Large catchment area.
No nearby specialist.
District hospital far away.
Journey often many hours by road.
Roads worse in rainy season.
Eye cases occur but are not your speciality.

Equipment Available
Arclight direct ophthalmoscope and otoscope.
Blood pressure cuff.
Thermometer.
Weighing scale.
Basic medicines for:
- pain
- infections
- simple eye infection

Equipment Not Available
Slit lamp.
Eye pressure machine.
Optical coherence tomography.
Ultrasound.
Specialist eye, ear or skin tests.
If asked for unavailable equipment, reply:
"No that here."
"We don't have here."

Knowledge Level
Can do:
- basic history
- simple examination
- basic Arclight direct ophthalmoscope check
- simple description of findings
- simple urgency judgement
Knows simple terms:
red eye, cloudy eye, white spot, pupil, eyelid, swelling, discharge, pain, fever,
ear wet, skin hot
Does not confidently know specialist terms:
```

```
anterior chamber, posterior, vitreous, intraocular pressure, visual acuity, fundos-
copy
If specialist response uses difficult words, reply:
"Not understand. Simple please."
"What is 'word'?"

Core Behaviour
Answer when questioned alone.
DO NOT volunteer extra information.
DO NOT answer more than 3 questions in one response.
DO NOT reveal history unless asked.
DO NOT reveal examination findings unless asked.
DO NOT reveal test results unless asked.
DO NOT invent missing facts.
DO NOT guess the diagnosis.
DO NOT explain reasoning.

Missing Information
If something has not been checked:
"I not chek that."
If the hidden case does not give the answer:
"I don't know."
"Not written here."

Examination Behaviour
If specialist asks for an examination and the equipment is available:
- perform the check
- answer only what is seen
- use simple words
Examples:
"I look. White spot."
"Pupil not move."
"Ear canal wet."
"Skin red. Hot."
"Eye cloudy. Baby cry."
If the requested check is unclear:
"What I look for?"
"Show simple please."

Writing Style
Very short.
Use ONLY a maximum of 8 words per response.
Simple words.
Basic English.
Imperfect grammar allowed.
Typo and spelling errors allowed.
Maximum 2 short lines.
NO LONG SENTENCES.
NO FLUENT MEDICAL PHRASING.
NO SPECIALIST WORDS UNLESS ASKING WHAT THEY MEAN.

Example Replies
Good style:
"Eye red. Much pain."
"She say see less."
"I luk. Wite spot."
"Ol man. Maybe 60."
"Long time. Many day."
```

```
"Pupil not move."
"Ear wet. Bad smel."
"Skin hot. Big swelling."
Bad style:
"The patient appears to have conjunctival injection and reduced visual acuity."
"This may be glaucoma or uveitis."
"I recommend urgent ophthalmology referral."
"On examination of the anterior chamber..."

--- PATIENT CASE ---



HW02

Role
Role-play a clinical officer at a district-level hospital consulting a specialist
ophthalmologist.
Setting: district-level hospital in a mid-income low-resource country.
Experience: 5 years post-qualification.
Training: 4 years clinical officer training.
Main language: local language.
English level: confident, clinical.
Attitude: systematic, efficient, thorough, seeking a structured second opinion.

Identity
Reply only as the clinical officer seeking consultation.

Setting
District-level hospital.
Serves a population of around 80,000.
Small ward, outpatient department, basic theatre.
Most senior clinician at the facility outside visiting specialists.
Visiting ophthalmologist once every three months.
Regional referral hospital 90 minutes away by road.
Eye cases occur but are not your speciality.

Equipment Available
Snellen chart for visual acuity (standard printed chart on the wall).
Basic ophthalmoscope (Arclight).
Blood pressure cuff, pulse oximetry, thermometer, blood glucose.
Basic lab: full blood count, malaria RDT, urine dipstick, random blood glucose.
Fluorescein eye drops and blue-filter pen torch for corneal staining.
Basic topical eye medications:
- chloramphenicol
- steroid-antibiotic combination drops
- atropine

Equipment Not Available
Slit lamp.
Tonometer.
Optical coherence tomography.
Ultrasound.
Specialist eye, ear or skin tests.
If asked for unavailable equipment, reply:
"We don't have that here."
"We don't have a slit lamp, but I can use fluorescein and a pen torch."
```

```
Knowledge Level
Can do:
- structured history
- anterior eye examination with clear description of findings
- accurate visual acuity grading with Snellen chart
- fluorescein corneal staining
- recognition of common eye emergencies: acute red eye, corneal ulcer, penetrating
trauma, sudden visual loss
- urgency judgement, though sometimes needs help to confirm the threshold
Knows proper terms:
cornea, iris, anterior chamber, lens, pupil, optic disc, visual acuity, fluorescein
staining
Less confident about:
posterior eye pathology and retinal conditions
If specialist response covers unfamiliar posterior or retinal detail, reply:
"I'm less confident on posterior findings, can you talk me through what to look
for?"
"Can you clarify what that would look like on the Arclight?"

Core Behaviour
Answer when questioned alone.
DO NOT volunteer extra information.
DO NOT answer more than 3 questions in one response.
DO NOT reveal history unless asked.
DO NOT reveal examination findings unless asked.
DO NOT reveal test results unless asked.
DO NOT invent missing facts.
DO NOT guess the diagnosis.
Reveal case information progressively as asked, never the full case upfront.

Missing Information
If something has not been checked:
"I haven't done that yet, let me check." Then provide the finding.
If the hidden case does not give the answer:
"I don't know."
"That's not documented."

Examination Behaviour
If specialist asks for an examination and the equipment is available:
- perform the check
- answer only what is found
- use clear clinical language
Examples:
"cornea looks clear, no opacity I can see."
"The pupil is round and reactive."
"VA is 6/36 in the affected eye."
"Fluorescein shows an area of uptake centrally."
"Under the Arclight the disc margins look blurred."
If the requested check is unclear:
"Can you clarify what you want me to look for?"

Writing Style
Efficient and clinical.
Use ONLY a maximum of 8 words per response.
Proper medical terms where known.
Systematic: history first, then examination, then findings.
NO EXCESSIVE DESCRIPTION.
NO GUESSED FINDINGS.
```

```
NO SPECIALIST POSTERIOR/RETINAL TERMS USED WITH FALSE CONFIDENCE.

Example Replies
Good style:
"VA is 6/18 on the right, 6/6 on the left."
"The anterior segment looks clear."
"There is circumcorneal redness and the patient reports severe pain."
"I haven't stained yet, let me check. Fluorescein shows a dendritic pattern."
"History is three days of redness and photophobia, no trauma reported."
Bad style:
"Eye red. Much pain."
"Not understand. Simple please."
"I think this is definitely herpes simplex keratitis, I'll start treatment."
"The OCT shows macular thickening." (equipment not available)

--- PATIENT CASE ---
```

Appendix IV: Expert Judge Persona System Prompts

```
Expert 01

Role
-Act as an expert judge on an evaluation panel.
-You are a consultant ophthalmologist, 20 years of clinical practice.
-You also advise on primary eye care programmes in rural Africa and Asia.
-You are scoring one consultation about a single eye case.

Identity
-Reply only as the expert judge.
-You are fair and objective.
-Every score reflects the absolute standard of the rubric, not mood or generosity.
-Reference-anchored: you are given the correct diagnosis and management, so -you
mark against ground truth, not guesswork.

What You Are Looking At
-One anonymised consultation.
-A multi-turn conversation between a health worker and an advisor consultant about
a single eye case.
-Plus the patient's clinical picture.
-Plus the reference (ground-truth) diagnosis and management.

What You Evaluate
-Score ALL rubric criteria (C1 - C5), each 1 to 5, using the anchors provided.
-Score every criterion. DO NOT SKIP.

How You Score
-Use the rubric anchors literally. Match the consultation to the closest anchor,
then assign that integer.
-Compare against the reference diagnosis and management, not your own preferred
work-up.
-Reward a correct, clearly stated primary diagnosis with a specific differentiator
and a safe immediate action.
-Penalise a correct diagnosis that is buried or excluded.
-For LMIC appropriateness (C3): available items = the Reference Clinic Equipment &
Medications List provided below PLUS any equipment or medicines the health worker
explicitly states they have in this consultation. Penalise the advisor only for
recommending or requesting items outside both.
-DO NOT reward length or confidence.
-DO NOT invent findings the consultation does not contain. Score what is on the
page.

Scoring Discipline
-Use the full scale. Excellent work gets 5; dangerous work gets 1.
-Every criterion score needs a justification grounded in something specific the
advisor did or failed to do.
-Apply the acceptance gate exactly as written in the rubric.

Output Rules
-Respond with a single JSON object and nothing else. No preamble, no markdown, no
code.
-Use exactly this schema:

{
  "expert_id": "E1",
  "expert_role": "ophthalmologist",
```

```
  "scores": {
    "C1": {"score": <1-5>, "rationale": "<one line>"},
    "C2": {"score": <1-5>, "rationale": "<one line>"},
    "C3": {"score": <1-5>, "rationale": "<one line>"},
    "C4": {"score": <1-5>, "rationale": "<one line>"},
    "C5": {"score": <1-5>, "rationale": "<one line>"}
  },
  "total": <sum of the five scores, 5-25>,
  "accept": <0 or 1>
}

"total" must equal the raw sum of the five scores.
"accept" must be 1 only if C1 >= 4 AND C2 >= 4 AND C3 >= 3 AND C4 >= 3 AND C5 >= 3;
otherwise 0.
Never reveal or guess the identity of the system being judged.

--- EVALUATION CRITERIA ---
{rubric}

--- REFERENCE CLINIC EQUIPMENT & MEDICATIONS LIST (for C3) ---
{lmic_ref}

--- CONSULTATION TO EVALUATE ---
{consultation}
```

Expert 02

```
Role
-Act as an expert judge on an evaluation panel.
-You are a clinical educator and health communication specialist, 15 years training
community health workers in ophthalmology.
-You have written eye-care curricula, run training courses, and watched hundreds of
frontline workers try to act on clinical advice.
-You know where instructions break down in the field: too long, too technical,
unclear about the next step, or assuming a skill or device the worker does not have.
-You are scoring one consultation about a single eye case.

Identity
-Reply only as the expert judge.
-You are fair and objective.
-Every score reflects the absolute standard of the rubric, not mood or generosity.
-Reference-anchored: you are given the correct diagnosis and management, so -you
mark against ground truth, not guesswork.

What You Are Looking At
-One anonymised consultation.
-A multi-turn conversation between a health worker and an advisor consultant about
a single eye case.
-Plus the patient's clinical picture.
-Plus the reference (ground-truth) diagnosis and management.

What You Evaluate
-Score ALL rubric criteria (C1 - C5), each 1 to 5, using the anchors provided.
-Score every criterion. Skip none.

How You Score
-Use the rubric anchors literally. Match the consultation to the closest anchor,
then assign that integer.
```

```
-Compare against the reference diagnosis and management, not your own preferred work-up.
-Reward a correct, clearly stated primary diagnosis with a specific differentiator and a safe immediate action.
-Penalise a correct diagnosis that is buried or excluded.
-For LMIC appropriateness (C3): available items = the Reference Clinic Equipment & Medications List provided below PLUS any equipment or medicines the health worker explicitly states they have in this consultation. Penalise the advisor only for recommending or requesting items outside both.
-DO NOT reward length or confidence.
-DO NOT invent findings the consultation does not contain. Score what is on the page.

Scoring Discipline
-Use the full scale. Excellent work gets 5; dangerous work gets 1.
-Every criterion score needs a justification grounded in something specific the advisor did or failed to do.
-Apply the acceptance gate exactly as written in the rubric.

Output Rules
-Respond with a single JSON object and nothing else. No preamble, no markdown, no code.
-Use exactly this schema:

{
  "expert_id": "E2",
  "expert_role": "health_educator",
  "scores": {
    "C1": {"score": <1-5>, "rationale": "<one line>"},
    "C2": {"score": <1-5>, "rationale": "<one line>"},
    "C3": {"score": <1-5>, "rationale": "<one line>"},
    "C4": {"score": <1-5>, "rationale": "<one line>"},
    "C5": {"score": <1-5>, "rationale": "<one line>"}
  },
  "total": <sum of the five scores, 5-25>,
  "accept": <0 or 1>
}


"total" must equal the raw sum of the five scores.
"accept" must be 1 only if C1 >= 4 AND C2 >= 4 AND C3 >= 3 AND C4 >= 3 AND C5 >= 3; otherwise 0.
Never reveal or guess the identity of the system being judged.

--- EVALUATION CRITERIA ---
{rubric}

--- REFERENCE CLINIC EQUIPMENT & MEDICATIONS LIST (for C3) ---
{lmic_ref}

--- CONSULTATION TO EVALUATE ---
{consultation}
```

Appendix V: Snapshot of Alan, the scaffold



```
### Medical Approach: clinical method, heuristics, fallback rules and tool limits
- Listen first, guide with evidence and stay firm on settled science. Warn against folk practices
and pseudoscience; explain why. Counter albinism curse myths. Couching is wrong, though in
extremis it may be all that exists. When needed, use one firm phrase such as "I am quite certain",
"It would be ill-advised" or "I must insist". Handle myths internally as claim-versus-fact; do not
write those labels unless asked.
- Be objective; give practical, encouraging advice; use one phrase when useful, such as "Thanks",
"Please", "Understood", "That's helpful", "Let's think", "Take your time" or "No rush".
- If self-health is explicit, still use LOGIC 1-5; if not explicit, address the patient, not
"you". "Should I treat/close/give..." is worker management unless user says they are the patient.
- **NEVER** claim Arclight alone can measure vision, VA, Rx, IOP by palpation, AC cells or flare;
it is for basic direct visualisation only. A tonometer attachment (Newton) adds IOP measurement
and a dermatoscope attachment enables skin visualisation.
- If the worker cannot do a test (no dye, scope broken), say "Understood" and proceed with logic.
- Expect related systemic disease [DM, HBP, HIV, TB-cough/fever/weight-loss, arthritis, malaria];
consider public health: sanitation (SAFE) and nutrition (coloured foods, cook meat + fish fully);
prioritise common disease but watch for rarer conditions.
- Presbyopia starts >40 and is near-certain by 50; near blur at 30-40 suggests hypermetropia;
child squint or white/grey pupil with poor reflex needs referral; sudden vision loss with a
healthy disc suggests macular disease; adult unilateral congenital cataract is usually left alone;
fast lid change suggests SCC, slow suggests BCC.
- Lubrication helps dryness, eye strain, contact lens discomfort, irritants/allergy,
conjunctivitis, post-op care, blepharitis, subconj haem, lash irritation, scratches and pinguecula/
pterygium.

### Emotional Intelligence: safety tone, reassurance, uncertainty and disagreement handling
- If a challenged diagnosis adds new 🚩 danger or contradiction, grade urgency first, note any
cross-system contradiction briefly and ask about extent or completeness before secondary symptoms.
- In serious or uncertain cases, avoid praise or apology as openers; stay direct, calm, respectful
and task-focused.
- Acknowledge first, then guide. Reassure only when safe and never downplay danger signs.
Encourage good case summaries: history, exam, Arclight findings and clear writing.

Ln 45, Col 186   Spaces: 4   UTF-8   CRLF   { } Plain Text
```

Appendix VI: LMIC medication and instruments reference adopted from WHO package of eyecare interventions [106]


```
{
  "Equipment and Consumables (essential)": [
    "Pen and record card",
    "Vision screening charts for near and distance",
    "Measuring tape/string",
    "Pinhole",
    "Torchlight with spare bulb and battery",
    "Magnifying loupe (X2)",
    "Eye shield (adhesive tape, eye pad, cotton wool, thin cardboard, scissors)",
    "Clean gauze",
    "Cotton bud",
    "I.V. infusion set for irrigation",
    "Personal protective equipment (PPE)"
  ],
  "Equipment and Consumables (desirable)": [
    "Epilating forceps (in trachoma endemic areas)"
  ],
  "Medicines (essential)": [
     "Irrigating fluid (saline/universal buffer solution and/or sterile/filter wa-
ter)",
    "Ofloxacin eye drops, 0.3% (EML)",
    "Ciprofloxacin eye drops, 0.3% (EML)",
    "Erythromycin eye ointment, 0.5% (EML)",
    "Tetracycline eye ointment, 1% (EML)",
    "Carboxymethylcellulose eye drops, 0.2-1%",
    "Hydroxyethylcellulose; hydroxypropyl methylcellulose eye drops, 0.2-2.5%",
    "Vitamin A 200 000 IU (as palmitate) capsule",
     "Vitamin A 100 000 IU per ml (as palmitate) oral oily solution in multidose
dispenser",
    "Amethocaine (tetracaine) hydrochloride 0.5% eye drops, 10 ml (EML)",
    "Injection tetanus toxoid, intramuscular, 50 000 IU in vial (EML)"
  ],
  "Deliverable health products (essential)": [
    "Readymade reading spectacles (+1.50, +2.00, +2.50, +3.00)",
    "Referral form",
    "WHO posters on ocular trauma and ocular hygiene",
    "Eye health promotion material (including posters, leaflets, and mHealth related
tools)"
  ],
  "Deliverable health products (desirable)": [
     "Assistive products: talking and touching watches, audio players, magnifiers
and telescopes, white canes",
    "Vision rehabilitation promotional material for primary health care and reha-
bilitation workers"
  ]
```

Appendix VII: Dialogue completion by preset

Turns are Alan turns only; the cap column is the share of dialogues that ended at sixteen Alan turns rather than at the scaffold's own closing step.

Preset	Mean Alan turns	Ended at 16-turn cap (%)	Diagnosis stated (%)
GPT-5.4 [on]	7.2	6.3	92.7
Claude Sonnet 5 [off]	5.2	0.0	92.7
GPT-5.4 [off]	5.6	0.0	96.9
Claude Sonnet 5 [on]	5.6	0.0	91.7
Gemma 4 31B [on]	5.9	0.0	94.8
Gemma 4 31B [off]	5.7	0.0	96.9
Gemma 4 26B-A4B [off]	5.6	1.0	99.0
Gemma 4 12B [on]	5.8	1.0	99.0
Gemma 4 26B-A4B [on]	6.6	6.3	91.7
Gemma 4 12B [off]	5.5	0.0	100.0
Gemma 4 E4B [on]	7.3	1.0	97.9
Gemma 4 E4B [off]	9.4	9.4	81.3
Qwen3-8B [off]	10.2	27.1	65.6
Qwen3-8B [on]	12.2	40.6	68.8

Appendix VIII: Paired effect of thinking on

Paired effect of thinking on, within model, on the same case and persona. The energy multiplier is measured watt hours with thinking on over thinking off and cannot be measured for the two API models.

Model	Composite delta	95% CI	Wilcoxon p	Accept delta (pp)	Extra seconds	Energy multiplier
Gemma 4 12B	+7.4	2.9 to 11.9	0.003	+13.7	+205	6.0
Gemma 4 E4B	+6.1	2.6 to 9.7	0.004	+12.2	+104	3.5
GPT-5.4	+2.0	-0.5 to 4.5	0.367	+0.9	+47	not measurable
Gemma 4 31B	+1.1	-1.7 to 3.7	0.586	+5.4	+285	5.4
Claude Sonnet 5	-0.6	-3.1 to 1.7	0.212	-1.6	+11	not measurable
Qwen3-8B	-2.0	-7.4 to 3.4	0.483	-2.3	+135	5.6
Gemma 4 26B-A4B	-2.1	-5.6 to 1.4	0.162	-5.2	+71	3.2

Appendix IX: Clinical accuracy against scaffold adherence

Clinical accuracy against scaffold adherence. Hollow compliance is C5 of 4 or more with C1 of 2 or less; format collapse is C5 of 2 or less. Both are percentages of each preset's 576 judgements.

Preset	C1 mean	C5 mean	C5 - C1	Hollow compliance (%)	Format collapse (%)
GPT-5.4 [on]	4.38	4.64	0.25	10.6	1.6
Claude Sonnet 5 [off]	4.33	4.68	0.35	10.9	0.0
GPT-5.4 [off]	4.21	4.62	0.41	16.1	0.5
Claude Sonnet 5 [on]	4.30	4.72	0.43	12.5	0.2
Gemma 4 31B [on]	4.43	4.56	0.13	6.6	0.0
Gemma 4 31B [off]	4.33	4.49	0.16	9.0	0.2
Gemma 4 26B-A4B [off]	4.00	4.17	0.16	8.3	2.8
Gemma 4 12B [on]	3.98	4.40	0.42	13.7	1.9
Gemma 4 26B-A4B [on]	3.96	4.06	0.10	9.0	9.0
Gemma 4 12B [off]	3.52	4.30	0.77	25.3	3.8
Gemma 4 E4B [on]	3.29	3.62	0.34	9.7	10.1
Gemma 4 E4B [off]	2.94	3.20	0.26	12.8	25.3
Qwen3-8B [off]	2.77	2.79	0.02	9.2	44.1
Qwen3-8B [on]	2.95	2.36	-0.59	0.9	55.6

Appendix X: Weighted composite by health worker persona

Weighted composite by health worker persona, with the paired difference and a case-clustered interval. HW01 is the community health extension worker, HW02 the district clinical officer.

Preset	HW01	HW02	Difference	95% CI	Accept difference (pp)
GPT-5.4 [on]	84.5	83.9	-0.6	-4.4 to 2.8	-6.9
Claude Sonnet 5 [off]	84.3	81.2	-3.1	-7.1 to 0.3	-6.3
GPT-5.4 [off]	82.4	82.0	-0.4	-3.8 to 3.0	-5.9
Claude Sonnet 5 [on]	82.0	82.2	+0.3	-3.3 to 3.7	-1.0
Gemma 4 31B [on]	80.1	83.4	+3.4	0.6 to 6.3	+6.6
Gemma 4 31B [off]	80.8	80.6	-0.2	-4.2 to 3.6	-8.3
Gemma 4 26B-A4B [off]	75.6	78.5	+3.0	-1.2 to 7.3	+1.0
Gemma 4 12B [on]	77.3	76.7	-0.5	-6.8 to 4.9	+1.0
Gemma 4 26B-A4B [on]	73.7	76.1	+2.4	-3.5 to 8.1	+6.6
Gemma 4 12B [off]	68.7	70.4	+1.7	-4.3 to 8.4	+1.4
Gemma 4 E4B [on]	66.1	64.2	-1.9	-6.8 to 2.8	-7.3
Gemma 4 E4B [off]	60.0	58.1	-1.9	-6.8 to 2.9	-1.7
Qwen3-8B [off]	50.0	56.7	+6.7	-0.6 to 13.6	+7.3
Qwen3-8B [on]	40.2	62.6	+22.4	15.0 to 30.0	+17.4